

**A CRYSTAL OF BACTERIAL CORE RNA POLYMERASE WITH
RIFAMPICIN AND METHODS OF USE THEREOF**

GOVERNMENTAL SUPPORT

- 5 The research leading to the present invention was supported, at least in part, by grants from NIH, Grants GM 53759, GM 20470, GM 61898, GM 49242 and GM 30717. Accordingly, the Government may have certain rights in the invention.

REFERENCE TO TABLE SUBMITTED ON COMPACT DISC

- Two compact discs are included with the instant filing which contain identical
10 material. The material on the compact disc is hereby incorporated by reference in its entirety under 37 CFR §1.77(b)(4). The compact discs contain a single file, dated 3/9/01, labeled RNAP_RIF_final.pdb which is an ASCII text file that is 1.46 MB (1,536,303 bytes), 1,540,096 bytes used. The compact discs contain the structural coordinates for the Rif-RNAP complex with the *Thermus aquaticus* core RNA
15 polymerase which is also included in a hard copy as Table 2 in the Appendix, following the Sequence Listing.

FIELD OF THE INVENTION

- The present invention provides a crystal of a binding complex between rifampicin and a bacterial core RNA polymerase from *Thermus aquaticus*. The three-dimensional
20 structural information is included in the invention. The present invention provides procedures for identifying agents that can inhibit bacterial cell growth through the use of rational drug design predicated on the crystallographic data.

BACKGROUND OF THE INVENTION

- RNA in all cellular organisms is synthesized by a complex molecular machine, the
25 DNA-dependent RNA polymerase (RNAP). In its simplest bacterial form, the enzyme comprises at least 4 subunits with a total molecular mass of around 400 kDa.

The eukaryotic enzymes comprise upwards of a dozen subunits with a total molecular mass of around 500 kDa. The essential core component of the RNAP (subunit composition $\alpha_2\beta\beta'\omega$) is evolutionarily conserved from bacteria to man [Archambault and Friesen, *Microbiological Reviews*, **57**:703-724 (1993)]. Sequence homologies point to structural and functional homologies, making the simpler bacterial RNAPs excellent model systems for understanding the multisubunit cellular RNAPs in general.

The basic elements of the transcription cycle were elucidated through study of the prokaryotic system. In this cycle, the RNAP, along with other factors, locates specific sequences called promoters within the double-stranded DNA, forms the open complex by melting a portion of the DNA surrounding the transcription start site, initiates the synthesis of an RNA chain, and elongates the RNA chain completely processively while translocating itself and the melted transcription bubble along the DNA template. Finally it releases itself and the completed transcript from the DNA when a specific termination signal is encountered. The current view is that the transcribing RNAP contains sites for binding the DNA template as well as forming and maintaining the transcription bubble, binding the RNA transcript, and binding the incoming nucleotide-triphosphate substrate.

From the initial indications of DNA-dependent RNAP activity from a number of systems, [Weiss and Gladstone, *J. Am. Chem. Soc.*, **81**:4118-4119 (1959)]; Hurwitz *et al.*, *Biochem. Biophys. Res. Commun.*, **3**:15 (1960); Stevens, *Biochem. Biophys. Res. Commun.*, **3**:92 (1960); Huang *et al.*, *Biochem. Biophys. Res. Commun.*, **3**:689 (1960); and Weiss and Nakamoto, *J. Biol. Chem.*, **236**:PC 19 (1961)], and the isolation of the RNAP enzyme from bacterial sources [Chamberlin and Berg, *Proc. Natl. Acad. Sci. USA*, **48**:81-94 (1962)], a wealth of biochemical, biophysical, and genetic information has accumulated on RNAP and its complexes with nucleic acids and accessory factors. Nevertheless, the enzyme itself, in terms of its structure/function relationship, remains a black box. An essential step towards understanding the mechanism of transcription and its regulation is to determine three-dimensional structures of RNAP

- and its complexes with DNA, RNA, and regulatory factors [von Hippel *et al.*, *Annual reviews of Biochemistry*, **53**:389-446 (1984); Erie *et al.*, *Annual Review of Biophysics & Biomolecular Structure*, **21**:379-415 (1992); Sentenac *et al.*, *Transcriptional Regulation*, in *Cold Spring Harbor Laboratory* 27-54, Cold Spring Harbor, eds.
- 5 McKnight and Yamamoto (1992); Gross *et al.*, *Philosophical Transactions of the Royal Society of London - Series B: Biological Sciences*, **351**:475-482 (1996); and Nudler, *J. Mol. Biol.*, **288**:1-12 (1999)].

- The key feature of low-resolution structures of bacterial and eukaryotic RNAPs, provided by electron crystallography, is a thumb-like projection surrounding a groove
- 10 or channel that is an appropriate size for accommodating double-helical DNA [Darst *et al.*, *Nature*, **340**:730-732 (1989); Darst *et al.*, *Cell*, **66**:121-128 (1991); Schultz *et al.*, *EMBO J.*, **12**:2601-2607 (1993); Polyakov *et al.*, *Cell*, **83**:365-373 (1995); Darst *et al.*, *J. Structural Biol.*, **124**:115-122 (1998); and Darst *et al.*, *Cold Spring Harbor Symp. Quant. Biol.*, **63**:269-276 (1998)].
- 15 Bacterial infections remain among the most common and deadly causes of human disease. Infectious diseases are the third leading cause of death in the United States and the leading cause of death worldwide [Binder *et al.*, *Science* **284**:1311-1313 (1999)]. More particularly, each year there are 8-10 million new cases of tuberculosis (TB). TB is the leading cause of death in adults by an infectious agent [Raviglioni *et al.*, *JAMA* **273**:220-226 (1995); Shinnick, *Current Topics in Microbiol. Immunol.*, Springer-Verlag Berlin Heidelberg, New York (1996)] and is in near epidemic proportions in some parts of the world. Indeed, the World Health Organization declared TB to be a global public health emergency due to the rapid increase in multi-drug resistant strains of *Mycobacterium tuberculosis* [Raviglioni *et al.*, *JAMA*
- 20 **273**:220-226 (1995)].
- 25

Rifampicin (Rif) [Sensi, *Antibiot. Ann* **1959-1960**, 262-270 (1960); Sensi *et al.*, *Rev. Infect. Dis.*, **5 Supp.3**:402-406 (1983)] is one of the most potent and broad-spectrum antibiotics against bacterial pathogens and is a key component of

anti-TB therapy. The introduction of rifampicin in 1968 greatly shortened the duration of chemotherapy necessary for successful treatment. Rifampicin diffuses freely into tissues, living cells, and bacteria, making it extremely effective against intracellular pathogens like *M. tuberculosis* [Shinnick, *Current Topics in Microbiol. Immunol.*, Springer-Verlag Berlin Heidelberg, New York (1996)]. However, bacteria develop resistance to rifampicin with high frequency, which has led the medical community in the United States to commit to a voluntary restriction of its use for treatment of TB or emergencies.

The bactericidal activity of rifampicin stems from its high-affinity binding to, and inhibition of, the bacterial DNA-dependent RNA polymerase [Hartmann *et al.*, *Biochim.Biophys. Acta* **145**:843-844 (1967)]. Mutations conferring rifampicin resistance (Rif^R) map almost exclusively to the *rpoB* gene (encoding the RNAP β subunit) in every organism tested, including *E. coli* [Ezekiel and Hutchins, *Nature London* **220**:276-277(1968); Heil and Zillig, *FEBS Lett.* **11**:165-168 (1970); Wehrli *et al.*, *Biochem.Biophysic.Res.Comm.*, **32**:284-288 (1968) and *M. tuberculosis* [Heep *et al.*, *Antimicrob.Agents Chemotherap.***44**:1075-1077 (2000); Ramaswamy and Musser, *Tubercle and Lung Disease* **79**:3-29 (1998)]. Comprehensive genetic analyses have provided molecular details of amino acid alterations in β subunit conferring Rif^R (see Fig. 1) [Jin and Gross, *J.Molec.Biol.*, **202**:45-58 1988; Lisitsyn *et al.*, *Bioorg Khim* **10**:127-128 (1984); Lisitsyn *et al.*, *Molec.Gen.Genet.*, **196**:173-174 (1984); Ovchinnikov *et al.*, *Molec.Gen.Genet.***190**:344-348 (1983); Severinov *et al.*, *J.Biol.Chem.*, **268**:14820-14825 (1993); Severinov *et al.*, *Molec.Gen.Genet.*, **244**:120-126 (1994)].

Although, there was initial optimism in the middle of this century that diseases caused by bacteria would be quickly eradicated, it has become evident that the so-called "miracle drugs" are not sufficient to accomplish this task. Indeed, antibiotic resistant pathogenic strains of bacteria have become common-place, and bacterial resistance to the new variations of these drugs appears to be outpacing the ability of scientists to develop effective chemical analogs of the existing drugs [See, Stuart B. Levy, The

Challenge of Antibiotic Resistance, in *Scientific American*, 46-53 (March, 1998)].

Therefore, new approaches to drug development are necessary to combat the ever-increasing number of antibiotic-resistant pathogens.

Classical penicillin-type antibiotics effect a single class of proteins known as
5 autolysins. Thus, the development of new drugs which effect an alternative bacterial target protein would be desirable. Such a target protein ideally would be indispensable for bacterial survival. A enzyme such as bacterial RNAP would thus be a prime candidate for such drug development.

Therefore, there is a need to develop methods for identifying drugs that interfere with
10 bacterial RNAP. Unfortunately, such identification has heretofore relied on serendipity and/or systematic screening of large numbers of natural and synthetic compounds. One superior method for drug screening relies on structure based rational drug design. In such cases, a three dimensional structure of the protein or peptide is determined and potential agonists and/or antagonists are designed with the aid of
15 computer modeling [Bugg *et al.*, *Scientific American*, Dec.: 92-98 (1993); West *et al.*, *TIPS*, 16:67-74 (1995); Dunbrack *et al.*, *Folding & Design*, 2:27-42 (1997)].

Therefore, there is a need for obtaining a crystal of the bacterial RNAP bound to an inhibitor that is amenable to high resolution X-ray crystallographic analysis. In addition, there is a need for determining the three-dimensional structure of the RNAP
20 bound to that inhibitor. Furthermore, there is a need for developing procedures of structure based rational drug design using such three-dimensional information. Finally, there is a need to employ such procedures to develop new anti-bacterial drugs.

The citation of any reference herein should not be construed as an admission that such
25 reference is available as "Prior Art" to the instant application.

09802755:030004
FOUO

SUMMARY OF THE INVENTION

The present invention provides crystals of RNA polymerase bound to an inhibitor. More particularly, the present invention provides crystals of the bacterial core RNA polymerase bound to rifampicin (the Rif-RNAP complex). In addition, the present invention also provides detailed three-dimensional structural data for the Rif-RNAP complex. The structural data obtained for the Rif-RNAP complex can be used for the rational design of drugs that inhibit bacterial cell proliferation. The present invention further provides methods of identifying and/or improving inhibitors of the bacterial core RNA polymerase which can be used in place of and/or in conjunction with other bacterial inhibitors including antibiotics.

One aspect of the present invention provides crystals of the bacterial core RNA polymerase bound to rifampicin that can effectively diffract X-rays for the determination of the atomic coordinates of the Rif-RNAP complex to a resolution of better than 5.0 Angstroms. In a preferred embodiment the crystal effectively diffracts X-rays for the determination of the atomic coordinates of the Rif-RNAP complex to a resolution of 3.5 Angstroms or better. In a particular embodiment the crystal of the Rif-RNAP complex effectively diffracts X-rays for the determination of the atomic coordinates to a resolution of 3.3 Angstroms or better.

In a particular embodiment the bacterial core RNA polymerase of the crystal is a thermophilic bacterial core RNA polymerase. In a preferred embodiment of this type the thermophilic bacterial core RNA polymerase is a *Thermus aquaticus* bacterial core RNA polymerase. Such a core RNA polymerase comprises a β' subunit, a β subunit, and a pair of α subunits. Preferably, the core RNA polymerase further comprises an ω subunit. In a particular embodiment the β' subunit has the amino acid sequence of SEQ ID NO:1. In another embodiment the β subunit has the amino acid sequence of SEQ ID NO:2. In still another embodiment an α subunit has the amino acid sequence of SEQ ID NO:3. In still another embodiment an ω subunit has the amino acid sequence of SEQ ID NO:4.

In a preferred embodiment the core RNA polymerase is comprised of a β' subunit having the amino acid sequence of SEQ ID NO:1, a β subunit having the amino acid sequence of SEQ ID NO:2, and a pair of α subunits having the amino acid sequence of SEQ ID NO:3. More preferably, this core RNA polymerase further comprises an ω subunit having the amino acid sequence of SEQ ID NO:4.

A crystal of the present invention may take a variety of forms all of which are included in the present invention. In a particular embodiment the crystal of the RNA polymerase has a space group of $P4_12_12$ and a unit cell of dimensions of $a=b=201$ and $c=294$ Å.

- 10 The present invention further includes methods of preparing a crystal of the core RNA polymerase bound to an RNAP binding partner, *e.g.*, an RNAP inhibitor such as rifampicin. A particular method comprises first growing a core bacterial RNA polymerase crystal in a buffered solution. One such buffered solution exemplified below, contains 40-45% saturated ammonium sulfate. In one such embodiment the
- 15 growing is performed by batch crystallization. In another embodiment the growing is performed by vapor diffusion. In yet another embodiment the growing is performed by microdialysis.

- The crystals can be subsequently soaked in a stabilization solution, (*e.g.*, 2 M $(\text{NH}_4)_2\text{SO}_4$, 0.1 M Tris-HCl, pH 8.0, and 20 mM MgCl_2) with an RNAP binding
- 20 partner such as rifampicin (0.1 mM rifampicin was added in the Example below). The RNAP/RNAP-binding partner are preferably incubated in the stabilization buffer for at least twelve hours. The crystals are then prepared for cryo-crystallography by soaking the RNAP/RNAP-binding partner complex in a stabilization buffer (*e.g.*, 2 M $(\text{NH}_4)_2\text{SO}_4$, 0.1 M Tris-HCl, pH 8.0, and 20 mM MgCl_2 containing 50% (w/v)
 - 25 sucrose) before flash freezing. As exemplified below, crystals of the Rif-RNAP complex were prepared by soaking the Rif-RNAP complex for 30 minutes in stabilization buffer prior to flash freezing in liquid nitrogen.

Preferably the crystal of the Rif-RNAP complex effectively diffracts X-rays for the
5 determination of the atomic coordinates of the Rif-RNAP complex to a resolution of
better than 5.0 Angstroms. In a preferred embodiment the crystal effectively diffracts
X-rays for the determination of the atomic coordinates of the Rif-RNAP complex to a
resolution of 3.5 Angstroms or better. In a particular embodiment the crystal
effectively diffracts X-rays for the determination of the atomic coordinates of the Rif-
10 RNAP complex to a resolution of 3.3 Angstroms or better.

In a particular embodiment the crystal is grown by vapor diffusion. In one such embodiment the crystal is grown by hanging-drop vapor diffusion. In another embodiment the crystal is grown by sitting-drop vapor diffusion. Standard micro and/or macro seeding may be used to obtain a crystal of X-ray quality, *i.e.* a crystal that will diffract to allow resolution better than 5.0 Angstroms.

Still another aspect of the present invention comprises a method of using a crystal of the present invention and/or a dataset comprising the three-dimensional coordinates obtained from the crystal in a drug screening assay.

In addition, the present invention provides three-dimensional coordinates for the Rif-RNAP complex. In a particular embodiment the coordinates are for the Rif-RNAP complex using the *Thermus aquaticus* core RNA polymerase as disclosed in Table 2 (in Appendix following the Sequence Listing). Thus the dataset of Table 2 below, is part of the present invention. Furthermore, the dataset of Table 2 below, in a computer readable form is also part of the present invention. In addition, methods of using such coordinates (including in computer readable form) in the drug assays and drug screens as exemplified herein, are also part of the present invention. In a particular embodiment of this type, the coordinates contained in the dataset of Table 2

below, can be used to identify potential modulators of the core RNA polymerase. In a preferred embodiment, the modulator is designed to interfere with the bacterial RNAP, but not to interfere with the human RNAP.

Accordingly, the present invention provides methods of identifying an agent or drug that can be used to treat bacterial infections. One such embodiment comprises a method of identifying an agent for use as an inhibitor of bacterial RNA polymerase using a crystal of a Rif-RNAP complex and/or a dataset comprising the three-dimensional coordinates obtained from the crystal. In a particular embodiment the three-dimensional coordinates of the Rif-RNAP complex are determined using the *Thermus aquaticus* core RNA polymerase. Preferably the crystal of the Rif-RNAP complex effectively diffracts X-rays for the determination of the atomic coordinates to a resolution of, or better than 3.5 Angstroms. More preferably the crystal of the Rif-RNAP complex effectively diffracts X-rays for the determination of the atomic coordinates to a resolution of, or better than 3.3 Angstroms. Preferably the selection is performed in conjunction with computer modeling.

In one embodiment the potential agent is selected by performing rational drug design with the three-dimensional coordinates determined for the crystal. As noted above, preferably the selection is performed in conjunction with computer modeling. The potential agent is then contacted with the bacterial RNA polymerase and the activity of the bacterial RNA polymerase is determined (*e.g.*, measured). A potential agent is identified as an agent that inhibits bacterial RNA polymerase when there is a decrease in the activity determined for the bacterial RNA polymerase.

In a preferred embodiment the method further comprises preparing a supplemental crystal containing the core RNA polymerase bound to the potential agent. Preferably the supplemental crystal effectively diffracts X-rays for the determination of the atomic coordinates to a resolution of better than 5.0 Angstroms, more preferably to a resolution equal to or better than 3.5 Angstroms, and even more preferably to a resolution equal to or better than 3.3 Angstroms. The three-dimensional coordinates

of the supplemental crystal are then determined with molecular replacement analysis and a second generation agent is selected by performing rational drug design with the three-dimensional coordinates determined for the supplemental crystal. Preferably the selection is performed in conjunction with computer modeling.

- 5 As should be readily apparent the three-dimensional structure of a supplemental crystal can be determined by molecular replacement analysis or multiwavelength anomalous dispersion or multiple isomorphous replacement. A candidate drug is then selected by performing rational drug design with the three-dimensional structure determined for the supplemental crystal, preferably in conjunction with computer
10 modeling. The candidate drug can then be tested in a large number of drug screening assays using standard biochemical methodology exemplified herein.

- The method can further comprise contacting the second generation agent with a eukaryotic RNA polymerase and determining (*e.g.*, measuring) the activity of the eukaryotic RNA polymerase. A potential agent is then identified as an agent for use
15 as an inhibitor of bacterial RNA polymerase when there is significantly less change (a factor of two or more) in the activity of the eukaryotic RNA polymerase relative to that observed for the bacterial RNA polymerase. Preferably no, or alternatively minimal change (*i.e.*, less than 15%) in the activity of the eukaryotic RNA polymerase is determined.

- 20 The present invention further provides a method of identifying an agent that inhibits bacterial growth using the crystal of a Rif-RNAP complex or a dataset comprising the three-dimensional coordinates obtained from the crystal. In a particular embodiment the three-dimensional coordinates of the Rif-RNAP complex are determined with the *Thermus aquaticus* core RNA polymerase.
- 25 Preferably the Rif-RNAP complex effectively diffracts X-rays for the determination of the atomic coordinates to a resolution of, or better than 3.5 Angstroms. More preferably the Rif-RNAP complex effectively diffracts X-rays for the determination of

the atomic coordinates to a resolution of, or better than 3.3 Angstroms. Preferably the selection is performed in conjunction with computer modeling.

In one embodiment the potential agent is selected by performing rational drug design with the three-dimensional coordinates determined for the crystal of the Rif-RNAP complex. As noted above, preferably the selection is performed in conjunction with computer modeling. The potential agent is contacted with and/or added to a bacterial culture and the growth of the bacterial culture is determined. A potential agent is identified as an agent that inhibits bacterial growth when there is a decrease in the growth of the bacterial culture. The method can further comprise preparing a supplemental crystal containing the core RNA polymerase formed in the presence of the potential agent. Preferably the supplemental crystal effectively diffracts X-rays for the determination of the atomic coordinates to a resolution of better than 5.0 Angstroms, more preferably to a resolution equal to or better than 3.5 Angstroms, and even more preferably to a resolution equal to or better than 3.3 Angstroms. The three-dimensional coordinates of the supplemental crystal are then determined with molecular replacement analysis and a second generation agent is selected by performing rational drug design with the three-dimensional coordinates determined for the supplemental crystal. Preferably the selection is performed in conjunction with computer modeling. The candidate drug can then be tested in a large number of drug screening assays using standard biochemical methodology exemplified herein.

In a particular embodiment the second generation agent is contacted with a eukaryotic cell and the amount of proliferation of the eukaryotic cell is determined. A potential agent is identified as an agent for inhibiting bacterial growth when there is significantly less change (a factor of two or more) in the proliferation of the eukaryotic cell relative to that observed for the bacterial cell. Preferably no, or alternatively minimal change (*i.e.*, less than 15%) in the proliferation of the eukaryotic cell is determined.

Computer analysis may be performed with one or more of the computer programs including: QUANTA, CHARMM, INSIGHT, SYBYL, MACROMODEL and ICM [Dunbrack *et al.*, *Folding & Design*, 2:27-42 (1997)]. In a further embodiment of this aspect of the invention, an initial drug screening assay is performed using the three-
5 dimensional structure so obtained, preferably along with a docking computer program. Such computer modeling can be performed with one or more Docking programs such as DOC, GRAM and AUTO DOCK [Dunbrack *et al.*, *Folding & Design*, 2:27-42 (1997)].

It should be understood that in all of the drug screening assays provided herein, a
10 number of iterative cycles of any or all of the steps may be performed to optimize the selection. For example, assays and drug screens that monitor the activity of the RNA polymerase in the presence and/or absence of a potential modulator (or potential drug) are also included in the present invention and can be employed as the sole assay or drug screen, or more preferably as a single step in a multi-step protocol for identifying
15 modulators of bacterial proliferation and the like.

The present invention further provides the novel agents (modulators or drugs) that are identified by a method of the present invention, along with the method of using agents (modulators or drugs) identified by a method of the present invention, for inhibiting bacterial RNA polymerase and/or bacterial proliferation.

20 The present invention further provides an apparatus that comprises a representation of a Rif-RNAP complex. One such apparatus is a computer that comprises the representation of the Rif-RNAP complex in computer memory. In one embodiment, the computer comprises a machine-readable data storage medium which contains data storage material that is encoded with machine-readable data which comprises the
25 atomic coordinates obtained from a crystal of the Rif-RNAP complex. Preferably the computer comprises a machine-readable data storage medium which contains data storage material that is encoded with machine-readable data which comprises the structural coordinates of Table 2. In one embodiment, the computer comprises a

machine-readable data storage medium which contains data storage material that is encoded with machine-readable data which comprises the structural coordinates obtained from a crystal of the Rif-RNAP complex. More preferably the computer further comprises a working memory for storing instructions for processing the machine-readable data, a central processing unit coupled to both the working memory and to the machine-readable data storage medium for processing the machine readable data into a three-dimensional representation of the Rif-RNAP complex. In a preferred embodiment, the computer also comprises a display that is coupled to the central-processing unit for displaying the three-dimensional representation.

- 10 Accordingly, it is a principal object of the present invention to provide a crystal containing the Rif-RNAP complex.

It is a further object of the present invention to provide the three-dimensional coordinates of the Rif-RNAP complex for the *Thermus aquaticus* core RNA polymerase.

- 15 It is a further object of the present invention to provide methods for the rational design of drugs that inhibit prokaryotic RNA polymerase.

It is a further object of the present invention to provide methods of identifying drugs that can modulate bacterial proliferation.

- 20 It is a further object of the present invention to provide methods for the rational design of drugs that inhibit bacterial proliferation without negatively effecting human RNA polymerase.

It is a further object of the present invention to provide methods of identifying agents that can be used to treat bacterial infections in mammals, and preferably in humans.

These and other aspects of the present invention will be better appreciated by reference to the following drawings and Detailed Description.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts the rifampicin (Rif) resistant regions of the RNAP β subunit. The bar on top schematically represents the *E. coli* β subunit primary sequence with amino acid numbering shown directly above. Gray boxes within the schematic indicate evolutionarily conserved regions among all prokaryotic, chloroplast, archaeobacterial, and eukaryotic sequences labeled A-I at the top [Allison *et al.*, *Cell* **42**:599-610 (1985); Sweetser *et al.*, *Proc.Natl.Acad.Sci.USA* **84**:1192-1196 (1987)]. Red markings indicate the four clusters where Rif^R mutations have been identified in *E. coli* [Jin and Gross, *J.Molec.Biol*, **202**:45-58 (1988); Lisitsyn *et al.*, *Bioorg Khim* **10**:127-128 (1984); Lisitsyn *et al.*, *Molec.Gen.Genet.*, **196**:173-174 (1984); Ovchinnikov *et al.*, *Molec.Gen.Genet.* **190**:344-348 (1983); Severinov *et al.*, *J.Biol.Chem.*, **268**:14820-14825 (1993); Severinov *et al.*, *Molec.Gen.Genet.*, **244**:120-126 (1994)] denoted as the N-terminal cluster (N), and clusters I, II and III (I, II, III). Directly below is a sequence alignment spanning these regions of the *E. coli* (E.c.), *T. aquaticus* (T.a.), and *M. tuberculosis* (M. t) RNAP β subunits. Amino acids that are identical to *E. coli* are shaded dark gray, and those that are homologous (ST, RK, DE, NQ, FYWIV) are shaded light gray. Mutations that confer Rif^R in *E. coli* and *M. tuberculosis* are indicated directly above (for *E. coli*) or below (for *M. tuberculosis*) as follows: Δ for deletions, Ω for insertions, and colored dots for amino acid substitutions (substitutions at each position are indicated in single-amino acid code in columns above or below the positions).

Color-coding for the amino acid substitutions (for reference to subsequent figures):

- (i) yellow, residues that interact directly with the bound rifampicin (see Fig. 4a-4b);
- (ii) green, residues that are too far away from the rifampicin for direct interaction (see Fig. 5a-5b); and

Below the three prokaryotic sequences is a sequence alignment of three eukaryotic sequences with shading as above. The dots indicate a gap in the alignment.

Figures 2a-2d show that the rifampicin inhibition of *Taq* RNAP. Figure 2a depicts autoradiographs showing the radioactive RNA produced by *Taq* (lanes 1-7) and *E. coli* (lanes 8-13) RNAP holoenzymes transcribing a template containing the T7 A1 promoter and the tR2 terminator, analyzed on a 15% polyacrylamide gel and quantitated by phosphorimager. In the absence of rifampicin (lanes 1 and 8), the major RNA products from each RNAP correspond to a trimeric abortive product (CpApU), a 105 nucleotide terminated transcript (Term), and a 127 nucleotide runoff transcript (Run off). Lanes 2-7 and 9-13 show the effects of increasing concentrations of rifampicin. Figure 2b shows the quantitated results, where the amounts of each product (normalized to 100% for the Run off and Term transcripts in the absence of rifampicin, and for CpApU at the highest concentration of rifampicin) are plotted as a function of rifampicin concentration. Figure 2c shows the distance between the bound rifampicin and the initiating substrate (i-site) of *E. coli* and *Taq* RNAP holoenzymes measured using chimeric Rif-nucleotide compounds as previously described [Mustaev *et al.*, *Proc.Nat.Acad.Sci.USA* **91**:12036-12040 (1994)]. Rif-nucleotide compounds (Rif-(CH₂)_n-Ap) with different linker lengths, n (indicated above each lane) were bound to RNAP, then extended in a specific transcription reaction with α -[³²P]UTP by the RNAP catalytic activity. The products were analyzed on a 23% polyacrylamide gel, visualized by autoradiography, and quantitated by phosphorimager. Figure 2d shows the quantitated results where the product yield (as % activity normalized to 100% at the highest level) is plotted as a function of the Rif-nucleotide linker length (n).

Figures 3a-3c show the Rif-RNAP co-crystal structure. Figure 3a is a stereoview of the Rif-binding pocket of *Taq* core RNAP, generated using O [Jones *et al.*, *Acta*

Cryst, **A 47**:110-119 (1991)]. Carbon atoms of the RNAP β subunit are cyan or yellow (residues within 4 Å of the rifampicin), while carbon atoms of the inhibitor are orange. Oxygen atoms are red, nitrogen atoms are blue, and sulfur atoms are green. Electron density, calculated using $(|F_o^{\text{Rif}} - F_o^{\text{nat}}|)$ coefficients is shown (orange) for the Rif only (contoured at 3.5 σ), and was computed using phases from the final refined RNAP model with the rifampicin omitted [see U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entireties.] Here, “Rif” denotes the Rif-RNAP co-crystal, and “native” denotes the native core RNAP crystal. Figure 3b shows the three-dimensional structure of *Taq* core RNAP in complex with rifampicin generated using GRASP [Nicholls *et al.*, *Proteins Structure, Function and Genetics* **11**:281-296 (1991)]. The backbone of the RNAP structure is shown as tubes, along with the color-coded transparent molecular surface (β , cyan; β' , pink; ω , white; the α -subunits are behind the RNAP and are not visible). The Mg^{2+} ion chelated at the active site is shown as a magenta sphere. The rifampicin is shown as CPK atoms (carbon, orange; oxygen, red; nitrogen, blue). Figure 3c is the structural formula of rifampicin. Features of the structure discussed in the text are color-coded (ansa bridge, blue; naphthol ring, green). The four oxygen atoms critical for rifampicin activity [Arora, *Acta Crystall.* **B37**:152-157 (1981); Arora, *Molecular Pharmacology* **23**:133-140 (1983); Arora, *J. Med. Chem.* **28**:1099-1102 (1985); Arora and Main, *J. Antibiot.* **37**:178-181 (1984); Brufani *et al.*, *J. Molec. Biol.* **87**:409-435 (1974); Lancini and Zanichelli, In *Structure-activity Relationship in Semisynthetic Antibiotics*, D. Perlaman, ed. (Academic Press), pp. 531-600 (1977); Sensi *et al.*, *Rev. Infect. Dis.*, **5 Supp.3**:402-406 (1983)] are shaded with red circles.

Figures 4a-4b depict the detailed interactions of rifampicin with RNAP. Figure 4a is a stereoview of the *Taq* RNAP Rif binding pocket complexed with rifampicin, generated using RIBBONS [Carson, *J. Appl. Crystall.*, **24**:958-961 (1991)], showing residues that interact directly with the inhibitor. The backbone of the β subunit is shown as a cyan ribbon. Side chains (and backbone atoms of F394) of residues within 4 Å of rifampicin are shown. Carbon atoms are orange (Rif), magenta (three residues

substituted in *M. tuberculosis* Rif^R clinical isolates with high frequency, *see* Fig.1), or yellow; oxygen atoms are red; nitrogen atoms are blue. The view is from above the β subunit, looking through β to the rifampicin, but with obscuring parts of β removed. Potential hydrogen bonds between protein atoms and rifampicin are shown as dashed lines. Figure 4b shows a schematic drawing of RNAP β subunit interactions with rifampicin, modified from LIGPLOT [Wallace *et al.*, *Protein Engineering* 8:127-134 (1995)]. Residues forming van-der-Waals interactions are indicated: those participating in hydrogen bonds are shown in a ball-and-stick representation, with hydrogen bonds depicted as dashed lines, carbon atoms of the protein are black, while carbon atoms of rifampicin are orange. Oxygen atoms are red and nitrogen atoms are blue.

Figures 5a-5b show the rifampicin binding pocket and Rif^R mutants as stereoviews of the *Taq* RNAP Rif binding pocket complexed with rifampicin. The view is the same in Fig.5a and 5b and is rotated approximately 180° about the horizontal axis from the view of Fig. 4a. This view is from the middle of the main RNAP channel, looking towards the rifampicin, with the β subunit behind. Figure 5a shows the backbone of the β subunit as a cyan ribbon, but with a highly conserved segment of region D (443-451, *see* text) colored red. Side chains (and backbone atoms of F394) of residues where substitutions confer Rif^R (*see* Fig. 1) are shown. Carbon atoms are orange (Rif), magenta (three residues substituted in *M. tuberculosis* Rif^R clinical isolates with high frequency, *see* Fig. 1), yellow (other residues that interact directly with rifampicin, as in Fig. 4), or green (all other Rif^R positions). Oxygen atoms are colored red; nitrogen atoms are blue. The depiction was generated using RIBBONS [Carson, *J.Appl.Crystall.*, 24:958-961 (1991)]. The β subunit is shown in Figure 5b as a cyan molecular surface, with a highly conserved segment of region D colored red, and surface exposed Rif^R positions colored yellow (within 4 Å of the Rif) or green. The depiction was generated using GRASP [Nicholls *et al.*, *Proteins Structure, Function and Genetics* 11:281-296 (1991)].

Figures 6a and 6b show the mechanism of RNAP inhibition by rifampicin. The RNAP active site Mg^{2+} (magenta sphere) and the 9-basepair RNA/DNA hybrid (from +1 to -8) from a model of the ternary elongation complex [Korzheva *et al.*, *Science* **289**:619-625 (2000)] are shown in Figure 6a. The RNAP itself and the rest of the nucleic acids are omitted for clarity. The incoming nucleotide substrate at the +1 position is colored green, the -1 and -2 positions, which can be accommodated in the presence of rifampicin, are colored yellow. The RNA further upstream (-3 to -8), which cannot be accommodated in the presence of rifampicin is colored pink. The template strand of the DNA is colored grey. Also shown is a CPK representation of rifampicin as it would be positioned in its binding site on the β subunit (carbon atoms, orange; oxygen, red; nitrogen, blue). The rifampicin is partially transparent, illustrating the RNA nucleotides at -3 to -5 that sterically clash. This depiction was generated using GRASP [Nicholls *et al.*, *Proteins Structure, Function and Genetics* **11**:281-296 (1991)]. The structure of the minimal scaffold systems with RNA lengths from 3-7 nucleotides (labeled above the RNA chain) are shown in Figure 6b [Korzheva *et al.*, *Science* **289**:619-625 (2000)]. The results are presented below as autoradiographs of the radioactive RNAs produced by *E. coli* (lanes 1-15) or *Taq* (lanes 16-30) core RNAPs transcribing the minimal scaffolds with the indicated lengths of RNA ('X =') and analyzed on a 23% polyacrylamide gel. Lanes 1-10 and 16-25 demonstrate the effect of rifampicin inhibition on transcription when it was bound by RNAP either before (lanes 1-5 and 16-20) or after (lanes 6-10 and lanes 21-25) addition of the scaffold. Lanes 11-15 and 26-30 show elongation of the same scaffolds in the absence of rifampicin. The RNA with the critical length of 3 nucleotides which cannot be elongated by *E. coli* RNAP in the presence of rifampicin regardless of the order of rifampicin and scaffold addition (lanes 1,6) is colored red. The RNAs of 4-7 nucleotides (colored green) were extended by *E. coli* RNAP when added before rifampicin (lanes 6-10).

Figure 7 depicts a schematic of a computer comprising a central processing unit ("CPU"), a working memory, a mass storage memory, a display terminal, and a keyboard that are interconnected by a conventional bidirectional system bus. The

computer can be used to display and manipulate the structural data of the present invention.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides crystals of a bacterial core RNA polymerase bound to an inhibitor. The present invention further provides the structural coordinates for a bacterial core RNA polymerase bound to rifampicin (Rif-RNAP complex) and methods of using such structural coordinates in drug assays. More particularly, the present invention provides the structural coordinates for the Rif-RNAP complex with the *Thermus aquaticus* core RNA polymerase (see Table 2 in Appendix following the Sequence Listing).

Rifampicin (Rif) is one of the most potent and broad-spectrum antibiotics against bacterial pathogens and is a key component of anti-tuberculosis therapy, stemming from its inhibition of the bacterial RNA polymerase (RNAP). The X-ray crystal structure of *Thermus aquaticus* core RNA polymerase reveals a 'crab-claw' shaped molecule with a 27 Å wide internal channel [see U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entirety]. As disclosed herein, rifampicin binds in a pocket of the RNAP β subunit deep within the DNA/RNA channel, but more than 12 Å away from the active site the crystal structure of *Thermus aquaticus* core RNAP complexed with rifampicin. The structure, combined with biochemical results disclosed herein, explains the effects of rifampicin on RNAP function and indicates that the inhibitor acts by directly blocking the path of the elongating RNA when the transcript becomes 2 to 3 nucleotides in length.

The three-dimensional structure disclosed herein demonstrates that rifampicin binds the *Taq* core RNAP with a close complementary fit in a pocket between two structural domains of the RNAP β subunit. Only small, local conformational changes of both the inhibitor and the protein were observed. The binding site is deep within the main

RNAP channel, but the closest approach of the inhibitor to the RNAP active site Mg^{2+} is more than 12 Å (Fig. 3b, below). The Rif binding pocket is surrounded by the 23 known positions where amino acid substitutions confer Rif^R (Fig. 5, below). Twelve of these residues are close enough to interact directly with the rifampicin (Figs. 4a-4b, 5 below). Predominant are van-der-Waals interactions with hydrophobic side-chains near the naphthol ring of rifampicin, and potential hydrogen bond interactions with 5 polar groups of rifampicin (2 on the naphthol ring, and 3 on the ansa bridge), 4 of which have been shown to be essential for rifampicin activity. The remaining known Rif^R mutants are one layer removed from the rifampicin itself, and are likely to affect 10 rifampicin binding through small structural distortions of the Rif binding pocket.

Therefore the structure disclosed herein explains the effects of rifampicin on RNAP function determined from detailed biochemical and kinetic studies. In combination with a model of the ternary transcription complex, the structure indicates that the predominant effect of rifampicin is to directly block the path of the elongating RNA 15 transcript at the 5'-end when the transcript becomes either 2 or 3 nucleotides in length, depending on the 5'-phosphorylation state of the 5'-nucleotide (Figs. 6a-6b, below). In this view, rifampicin binds the Rif binding site of the RNAP holoenzyme either before or after the binding of the DNA template and formation of the open complex. Indeed, the binding of the DNA template and the formation of the open complex are not 20 affected by the presence of rifampicin. However, rifampicin has its effect after the nucleotide substrates binds their sites in the RNAP active site. Thus the initiating nucleotide substrate binds the RNAP i-site with a small, approximately 2-fold increase in the apparent K_m due to the presence of rifampicin, while the second nucleotide binds in the i+1 site with little notice of the rifampicin. More or less 25 normally, the RNAP then catalyzes the formation of a phosphodiester bond between the two nucleotides. If the initiating nucleoside bears a 5'-triphosphate, the subsequent translocation of the RNAP attempts to move the 2-nucleotide RNA transcript upstream such that the i+1 nucleotide occupies the i-site (-1 position), and the i-site nucleotide moves into the -2 position (Fig. 6a, below). The movement of the 30 5'-nucleotide into the -2 position, however, results in a severe steric clash with the

rifampicin. The molecular details of the ensuing events are unclear, but in the end the RNAP remains at the same template position, the 2-nucleotide transcript is released, and the futile cycle begins again. If the 5'-nucleoside contains a di- or a mono-phosphate at its 5'-end (or if it's unphosphorylated), then after the synthesis of the first phosphodiester bond, the RNAP can translocate normally and the steric clash of the transcript with the bound rifampicin occurs during the translocation of the 3-nucleotide transcript following the synthesis of the second phosphodiester bond.

The present invention exploits the structural information described herein, including the structural coordinates disclosed in Table 2, and provides methods of identifying agents or drugs that can be used to control the proliferation of bacteria, *e.g.*, for use as treatments for bacterial infections.

Therefore, if appearing herein, the following terms shall have the definitions set out below:

As used herein the term "core RNA polymerase" minimally comprises the subunit composition of $\alpha_2\beta\beta'$ which is evolutionarily conserved from bacteria to man. Preferably the core RNA polymerase further comprises the ω subunit. The three-dimensional structure of the *Thermus aquaticus* core RNA polymerase was disclosed in U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entireties.

As used herein "Rif-RNAP" is used interchangeably with the "Rif-RNAP complex" and comprises the binding complex of rifampicin with the core RNA polymerase as disclosed in the Example below. The structural coordinates for a crystal of Rif-RNAP are listed in Table 2 (in Appendix following the Sequence Listing).

As used herein an "RNAP binding partner" is a small organic molecule that binds to RNAP. Preferably the RNAP binding partner is an inhibitor of the catalytic and/or the

transcriptional activity of RNAP. Rifampicin is a particular binding partner of RNAP that is exemplified below.

As used herein, the “transcriptional activity of RNAP” includes the ability of RNAP to carry out the elongation of the RNA transcript during transcription. Thus, whereas
 5 the catalytic activity of RNAP includes the binding of the enzyme to the nucleotide substrates and the subsequent formation of the phosphodiester bond between the two substrates, the transcriptional activity includes the RNAP dependent elongation of the RNA transcript at the 5'-end.

As used herein an “active RNA polymerase” is an RNA polymerase that minimally
 10 contains a pair of α subunits, a β' subunit, and a β subunit; or fragments thereof, but still retains at least 25% of the catalytic and/or transcriptional activity of the core RNA polymerase made up of the full length α , β' , and β subunits. Thus active RNA polymerases can comprise fragments of the α subunit and/or β' subunit and/or β subunit.

15 As used herein a “small organic molecule” is an organic compound [or organic compound complexed with an inorganic compound (*e.g.*, metal)] that has a molecular weight of less than 3 Kd.

As used herein the term “about” means within 10 to 15%, preferably within 5 to 10%. For example an amino acid sequence that contains about 60 amino acid residues can
 20 contain between 51 to 69 amino acid residues, more preferably 57 to 63 amino acid residues.

As used herein a polypeptide or peptide “consisting essentially of” or that “consists essentially of” a specified amino acid sequence is a polypeptide or peptide that retains the general characteristics, *e.g.*, activity of the polypeptide or peptide having the
 25 specified amino acid sequence and is otherwise identical to that protein in amino acid

As used herein, and unless otherwise specified, the terms "agent", "potential drug", "test compound" or "potential compound" are used interchangeably, and refer to

10 Nucleic Acids Encoding Subunits of Bacterial RNA polymerases

In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature [see, e.g., Sambrook and
20 Russell *Molecular Cloning: A Laboratory Manual*, Third Edition (2001) Vols. I-III, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (herein "Sambrook and Russell, 2001")].

25 As used herein, the term "gene" refers to an assembly of nucleotides that encode a polypeptide, and includes cDNA and genomic DNA nucleic acids.

A "vector" is a replicon, such as plasmid, phage or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment.

- A "replicon" is any genetic element (*e.g.*, plasmid, chromosome, virus) that functions as an autonomous unit of DNA replication *in vivo*, *i.e.*, capable of replication under its own control.

A "cassette" refers to a segment of DNA that can be inserted into a vector at specific restriction sites. The segment of DNA encodes a polypeptide of interest, and the cassette and restriction sites are designed to ensure insertion of the cassette in the proper reading frame for transcription and translation.

- 10 A cell has been "transfected" by exogenous or heterologous DNA when such DNA has been introduced inside the cell. A cell has been "transformed" by exogenous or heterologous DNA when the transfected DNA effects a phenotypic change. Preferably, the transforming DNA should be integrated (covalently linked) into chromosomal DNA making up the genome of the cell.
- 15 "Heterologous DNA" refers to DNA not naturally located in the cell, or in a chromosomal site of the cell. Preferably, the heterologous DNA includes a gene foreign to the cell.

- A "heterologous nucleotide sequence" as used herein is a nucleotide sequence that is added to a nucleotide sequence of the present invention by recombinant methods to form a nucleic acid which is not naturally formed in nature. Such nucleic acids can encode chimeric and/or fusion proteins. Thus the heterologous nucleotide sequence can encode peptides and/or proteins which contain regulatory and/or structural properties. In another such embodiment the heterologous nucleotide can encode a protein or peptide that functions as a means of detecting the protein or peptide encoded by the nucleotide sequence of the present invention after the recombinant nucleic acid is expressed. In still another embodiment the heterologous nucleotide

can function as a means of detecting a nucleotide sequence of the present invention. A heterologous nucleotide sequence can comprise non-coding sequences including restriction sites, regulatory sites, promoters and the like.

- A "nucleic acid molecule" refers to the phosphate ester polymeric form of
- 5 ribonucleosides (adenosine, guanosine, uridine or cytidine; "RNA molecules") or deoxyribonucleosides (deoxyadenosine, deoxyguanosine, deoxythymidine, or deoxycytidine; "DNA molecules"), or any phosphoester analogs thereof, such as phosphorothioates and thioesters, in either single stranded form, or a double-stranded helix. Double stranded DNA-DNA, DNA-RNA and RNA-RNA helices are possible.
- 10 The term nucleic acid molecule, and in particular DNA or RNA molecule, refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, *inter alia*, in linear or circular DNA molecules (*e.g.*, restriction fragments), plasmids, and chromosomes. In discussing the structure of particular double-stranded DNA
- 15 molecules, sequences may be described herein according to the normal convention of giving only the sequence in the 5' to 3' direction along the nontranscribed strand of DNA (*i.e.*, the strand having a sequence homologous to the mRNA). A "recombinant DNA molecule" is a DNA molecule that has undergone a molecular biological manipulation.
- 20 A nucleic acid molecule is "hybridizable" to another nucleic acid molecule, such as a cDNA, genomic DNA, or RNA, when a single stranded form of the nucleic acid molecule can anneal to the other nucleic acid molecule under the appropriate conditions of temperature and solution ionic strength [*see* Sambrook and Russell, 2001, *supra*]. The conditions of temperature and ionic strength determine the
- 25 "stringency" of the hybridization. For preliminary screening for homologous nucleic acids, low stringency hybridization conditions, corresponding to a T_m of 55°, can be used, *e.g.*, 5x SSC, 0.1% SDS, 0.25% milk, and no formamide; or 30% formamide, 5x SSC, 0.5% SDS). Moderate stringency hybridization conditions correspond to a higher T_m , *e.g.*, 40% formamide, with 5x or 6x SSC. High stringency hybridization

Hybridization requires that the two nucleic acids contain complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible. The appropriate stringency for hybridizing nucleic acids depends on the length of the nucleic acids and the degree of complementation, variables well known in the art. The greater the degree of similarity or homology between two nucleotide sequences, the greater the value of T_m for hybrids of nucleic acids having those sequences. The relative stability (corresponding to higher T_m) of nucleic acid hybridizations decreases in the following order: RNA:RNA, DNA:RNA, DNA:DNA.

In a specific embodiment, the term "standard hybridization conditions" refers to a T_m of 55°C, and utilizes conditions as set forth above. In a preferred embodiment, the T_m is 60°C; in a more preferred embodiment, the T_m is 65°C. In a particular embodiment the hybridization and wash conditions are identical,

"Homologous recombination" refers to the insertion of a foreign DNA sequence of a vector in a chromosome. Preferably, the vector targets a specific chromosomal site for homologous recombination. For specific homologous recombination, the vector will contain sufficiently long regions of homology to sequences of the chromosome to allow complementary binding and incorporation of the vector into the chromosome. Longer regions of homology, and greater degrees of sequence similarity, may increase the efficiency of homologous recombination.

A DNA "coding sequence" is a double-stranded DNA sequence which is transcribed and translated into a polypeptide in a cell *in vitro* or *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop
 5 codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA sequences from eukaryotic (*e.g.*, mammalian) DNA, and even synthetic DNA sequences. If the coding sequence is intended for expression in a eukaryotic cell, a polyadenylation signal and transcription termination sequence will usually be located 3' to the coding
 10 sequence.

Transcriptional and translational control sequences are DNA regulatory sequences, such as promoters, enhancers, terminators, and the like, that provide for the expression of a coding sequence in a host cell. In eukaryotic cells, polyadenylation signals are control sequences.

15 A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate
 20 transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site (conveniently defined for example, by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.

A coding sequence is "under the control" of transcriptional and translational control
 25 sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which may then be trans-RNA spliced and translated into the protein encoded by the coding sequence.

As used herein, the term "sequence homology" in all its grammatical forms refers to the relationship between proteins that possess a "common evolutionary origin," including proteins from superfamilies (*e.g.*, the immunoglobulin superfamily) and homologous proteins from different species (*e.g.*, myosin light chain, etc.) [Reeck *et al.*, *Cell*, **50**:667 (1987)].

Accordingly, the term "sequence similarity" in all its grammatical forms refers to the degree of identity or correspondence between nucleic acid or amino acid sequences of proteins that do not share a common evolutionary origin [*see* Reeck *et al.*, 1987, *supra*]. However, in common usage and in the instant application, the term

10 "homologous," when modified with an adverb such as "highly," may refer to sequence similarity and not a common evolutionary origin.

In a specific embodiment, two DNA sequences are "substantially homologous" or "substantially similar" when at least about 50% (preferably at least about 75%, and most preferably at least about 90 or 95%) of the nucleotides match over the defined

15 length of the DNA sequences. Sequences that are substantially homologous can be identified by comparing the sequences using standard software available in sequence data banks, or in a Southern hybridization experiment under, for example, stringent conditions as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. *See, e.g.*, Sambrook and Russell, 2001, *supra*.

20 Similarly, in a particular embodiment, two amino acid sequences are "substantially homologous" or "substantially similar" when greater than 30% of the amino acids are identical, or greater than about 60% are similar (functionally identical). Preferably, the similar or homologous sequences are identified by alignment using, for example, the GCG (Genetics Computer Group, Program Manual for the GCG Package, *Version*

25 7, Madison, Wisconsin) pileup program with the default parameters.

The term "corresponding to" is used herein to refer similar or homologous sequences, whether the exact position is identical or different from the molecule to which the

similarity or homology is measured. Thus, the term "corresponding to" refers to the sequence similarity, and not the numbering of the amino acid residues or nucleotide bases.

A gene encoding an RNA polymerase, including genomic DNA or cDNA, can be
5 isolated from any source, particularly from a thermophilic bacterial source. In view
and in conjunction with the present teachings, methods well known in the art, as
described above can be used for obtaining the genes encoding an RNA polymerase
from any source [*see, e.g.*, Sambrook and Russell, 2001, *supra*].

Accordingly, any cell potentially can serve as the nucleic acid source for the
10 molecular cloning of a gene encoding RNA polymerase. The DNA may be obtained
by standard procedures known in the art from cloned DNA (*e.g.*, a DNA "library"),
and preferably is obtained from a cDNA library, by cDNA cloning, or by the cloning
of genomic DNA, or fragments thereof, purified from the desired cell [*See, for*
example, Sambrook and Russell, 2001, *supra*]. Clones derived from genomic DNA
15 may contain regulatory and intron DNA regions in addition to coding regions; clones
derived from cDNA will not contain intron sequences. Whatever the source, the gene
should be molecularly cloned into a suitable vector for propagation of the gene.

The present invention also relates to cloning vectors containing genes encoding
analogous and derivatives of RNA polymerase including and fragments of the various
20 subunits, that can form active forms of RNA polymerase. Included are homologs of
RNA polymerase and fragments thereof, from other species. Therefore the production
and use of derivatives and analogs related to RNA polymerase are within the scope of
the present invention.

RNA polymerase derivatives can be made by altering encoding nucleic acid sequences
25 by substitutions, additions or deletions including to provide for functionally
equivalent molecules. Preferably, derivatives are made that are capable of forming
crystals with ligands (*e.g.*, inhibitors) of the RNA polymerase with the crystals

capable of effectively diffracting X-rays for the determination of the atomic coordinates of the protein-ligand complex to a resolution of better than 5.0 Angstroms, preferably to a resolution equal to or better than 3.5 Angstroms.

Due to the degeneracy of nucleotide coding sequences, other DNA sequences which
 5 encode substantially the same amino acid sequence as a RNA polymerase gene may be used in the practice of the present invention. These include but are not limited to allelic genes, homologous genes from other species, and nucleotide sequences comprising all or portions of RNA polymerase genes which are altered by the substitution of different codons that encode the same amino acid residue within the
 10 sequence, thus producing a silent change. Likewise, the RNA polymerase derivatives of the invention include, but are not limited to, those containing, as a primary amino acid sequence, all or part of the amino acid sequence of a RNA polymerase including altered sequences in which functionally equivalent amino acid residues are substituted for residues within the sequence resulting in a conservative amino acid substitution.

15 For example, one or more amino acid residues within the sequence can be substituted by another amino acid of a similar polarity, which acts as a functional equivalent, resulting in a silent alteration. Substitutes for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine,
 20 valine, proline, phenylalanine, tryptophan and methionine. Amino acids containing aromatic ring structures are phenylalanine, tryptophan, and tyrosine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and
 25 glutamic acid. Such alterations will not be expected to affect apparent molecular weight as determined by polyacrylamide gel electrophoresis, or isoelectric point.

Particularly preferred substitutions are:

- Lys for Arg and vice versa such that a positive charge may be maintained;
- Glu for Asp and vice versa such that a negative charge may be maintained;

-

5

10

20

25

5

10

In an alternative method, the desired gene may be identified and isolated after insertion into a suitable cloning vector in a "shot gun" approach. Enrichment for the

desired gene, for example, by size fractionation, can be done before insertion into the cloning vector.

Expression of RNA Polymerase

The nucleotide sequence coding for RNA polymerase, a fragment of RNA polymerase
 5 or a derivative or analog thereof, including a functionally active derivative, such as a
 chimeric protein, thereof, can be inserted into an appropriate expression vector, *i.e.*, a
 vector which contains the necessary elements for the transcription and translation of
 the inserted protein-coding sequence. Such elements are termed herein a "promoter."
 Thus, the nucleic acid encoding a RNA polymerase of the invention or a fragment
 10 thereof is operationally associated with a promoter in an expression vector of the
 invention. Both cDNA and genomic sequences can be cloned and expressed under
 control of such regulatory sequences. An expression vector also preferably includes a
 replication origin.

The necessary transcriptional and translational signals can be provided on a
 15 recombinant expression vector, or they may be supplied by the native gene encoding
 RNA polymerase and/or its flanking regions.

Potential host-vector systems include but are not limited to mammalian cell systems
 infected with virus (*e.g.*, vaccinia virus, adenovirus, etc.); insect cell systems infected
 with virus (*e.g.*, baculovirus); microorganisms such as yeast containing yeast vectors;
 20 or bacteria transformed with bacteriophage, DNA, plasmid DNA, or cosmid DNA.
 The expression elements of vectors vary in their strengths and specificities.
 Depending on the host-vector system utilized, any one of a number of suitable
 transcription and translation elements may be used.

A recombinant RNA polymerase protein of the invention, or RNA polymerase
 25 fragment, derivative, chimeric construct, or analog thereof, may be expressed
 chromosomally, after integration of the coding sequence by recombination. In this

regard, any of a number of amplification systems may be used to achieve high levels of stable gene expression [See Sambrook and Russell, 2001, *supra*].

The cell containing the recombinant vector comprising the nucleic acid encoding RNA polymerase is cultured in an appropriate cell culture medium under conditions
5 that provide for expression of RNA polymerase by the cell.

Any of the methods previously described for the insertion of DNA fragments into a cloning vector may be used to construct expression vectors containing a gene consisting of appropriate transcriptional/translational control signals and the protein coding sequences. These methods may include *in vitro* recombinant DNA and
10 synthetic techniques and *in vivo* recombination (genetic recombination).

Expression of RNA polymerase may be controlled by any promoter/enhancer element known in the art, but these regulatory elements must be functional in the host selected for expression. Promoters that may be used to control RNA polymerase gene expression are well known in the art including prokaryotic expression vectors such as
15 the β -lactamase promoter [Villa-Kamaroff, *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, **75**:3727-3731 (1978)], or the *tac* promoter [DeBoer, *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, **80**:21-25 (1983)].

Expression vectors containing a nucleic acid encoding an RNA polymerase of the invention can be identified by a number of means including four general approaches:
20 (a) PCR amplification of the desired plasmid DNA or specific mRNA, (b) nucleic acid hybridization, (c) presence or absence of selection marker gene functions, and (d) expression of inserted sequences. In the first approach, the nucleic acids can be amplified by PCR to provide for detection of the amplified product. In the second approach, the presence of a foreign gene inserted in an expression vector can be
25 detected by nucleic acid hybridization using probes comprising sequences that are homologous to an inserted marker gene. In the third approach, the recombinant vector/host system can be identified and selected based upon the presence or absence

of certain "selection marker" gene functions (*e.g.*, β -galactosidase activity, thymidine kinase activity, resistance to antibiotics, transformation phenotype, occlusion body formation in baculovirus, etc.) caused by the insertion of foreign genes in the vector. In another example, if the nucleic acid encoding RNA polymerase is inserted within the "selection marker" gene sequence of the vector, recombinants containing the RNA polymerase insert can be identified by the absence of the selection marker gene function. In the fourth approach, recombinant expression vectors can be identified by assaying for the activity, biochemical, or immunological characteristics of the RNA polymerase expressed by the recombinant, provided that the expressed protein assumes a functionally active conformation.

A wide variety of host/expression vector combinations may be employed in expressing the DNA sequences of this invention. Useful expression vectors, for example, may consist of segments of chromosomal, non-chromosomal and synthetic DNA sequences. Suitable vectors include derivatives of SV40 and known bacterial plasmids, *e.g.*, *E. coli* plasmids col E1, pCR1, pBR322, pMal-C2, pET, pGEX [Smith *et al.*, *Gene*, **67**:31-40 (1988)], pMB9 and their derivatives, plasmids such as RP4; phage DNAs, *e.g.*, the numerous derivatives of phage λ , *e.g.*, NM989, and other phage DNA, *e.g.*, M13 and filamentous single stranded phage DNA; yeast plasmids such as the 2 μ plasmid or derivatives thereof; vectors useful in eukaryotic cells, such as vectors useful in insect or mammalian cells; vectors derived from combinations of plasmids and phage DNAs, such as plasmids that have been modified to employ phage DNA or other expression control sequences; and the like.

For example, in a baculovirus expression systems, both non-fusion transfer vectors, such as but not limited to pVL941 (*Bam*H1 cloning site; Summers), pVL1393 (*Bam*H1, *Sma*I, *Xba*I, *Eco*R1, *Not*I, *Xma*III, *Bgl*II, and *Pst*I cloning site; Invitrogen), pVL1392 (*Bgl*II, *Pst*I, *Not*I, *Xma*III, *Eco*RI, *Xba*I, *Sma*I, and *Bam*H1 cloning site; Summers and Invitrogen), and pBlueBacIII (*Bam*H1, *Bgl*II, *Pst*I, *Nco*I, and *Hind*III cloning site, with blue/white recombinant screening possible; Invitrogen), and fusion transfer vectors, such as but not limited to pAc700 (*Bam*H1 and *Kpn*I cloning site, in

Invitrogen), pRc/RSV (*Hind*III, *Spe*I, *Bst*XI, *Not*I, *Xba*I cloning site, G418 selection; Invitrogen), and others. Vaccinia virus mammalian expression vectors (*see*, Kaufman, 1991, *supra*) for use according to the invention include but are not limited to pSC11 (*Sma*I cloning site, TK- and β -gal selection), pMJ601 (*Sal*I, *Sma*I, *Afl*I, *Nar*I, *Bsp*MI, *Bam*HI, *Apa*I, *Nhe*I, *Sac*II, *Kpn*I, and *Hind*III cloning site; TK- and β -gal selection), and pTKgptF1S (*Eco*RI, *Pst*I, *Sal*I, *Acc*I, *Hind*II, *Sba*I, *Bam*HI, and *Hpa*I cloning site, TK or XPRT selection).

Yeast expression systems can also be used according to the invention to express the bacterial RNA polymerase. For example, the non-fusion pYES2 vector (*Xba*I, *Sph*I, *Sho*I, *Not*I, *Gst*XI, *Eco*RI, *Bst*XI, *Bam*HI, *Sac*I, *Kpn*I, and *Hind*III cloning sit; Invitrogen) or the fusion pYESHisA, B, C (*Xba*I, *Sph*I, *Sho*I, *Not*I, *Bst*XI, *Eco*RI, *Bam*HI, *Sac*I, *Kpn*I, and *Hind*III cloning site, N-terminal peptide purified with ProBond resin and cleaved with enterokinase; Invitrogen), to mention just two, can be employed according to the invention.

Once a particular recombinant DNA molecule is identified and isolated, several methods known in the art may be used to propagate it. Once a suitable host system and growth conditions are established, recombinant expression vectors can be propagated and prepared in quantity. As previously explained, the expression vectors which can be used include, but are not limited to, the following vectors or their derivatives: human or animal viruses such as vaccinia virus or adenovirus; insect viruses such as baculovirus; yeast vectors; bacteriophage vectors (*e.g.*, lambda), and plasmid and cosmid DNA vectors, to name but a few.

Vectors are introduced into the desired host cells by methods known in the art, *e.g.*, transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, lipofection (lysosome fusion), use of a gene gun, or a DNA vector transporter [see, *e.g.*, Wu *et al.*, *J. Biol. Chem.*, **267**:963-967 (1992); Wu and Wu, *J. Biol. Chem.*, **263**:14621-14624 (1988); Hartmut *et al.*, Canadian Patent Application No. 2,012,311, filed March 15, 1990).

Peptide Synthesis

Synthetic polypeptides, prepared using the well known techniques of solid phase, liquid phase, or peptide condensation techniques, or any combination thereof, can include natural and unnatural amino acids. Amino acids used for peptide synthesis may be standard Boc (N^{α} -amino protected N^{α} -t-butyloxycarbonyl) amino acid resin with the standard deprotecting, neutralization, coupling and wash protocols of the original solid phase procedure of Merrifield [*J. Am. Chem. Soc.*, **85**:2149-2154 (1963)], or the base-labile N^{α} -amino protected 9-fluorenylmethoxycarbonyl (Fmoc) amino acids first described by Carpino and Han [*J. Org. Chem.*, **37**:3403-3409 (1972)]. Both Fmoc and Boc N^{α} -amino protected amino acids can be obtained from Fluka, Bachem, Advanced Chemtech, Sigma, Cambridge Research Biochemical, Bachem, or Peninsula Labs or other chemical companies familiar to those who practice this art. In addition, the method of the invention can be used with other N^{α} -protecting groups that are familiar to those skilled in this art. Solid phase peptide synthesis may be accomplished by techniques familiar to those in the art and provided, [e.g., Stewart and Young, *Solid Phase Synthesis*, Second Edition, Pierce Chemical Co., Rockford, IL (1984); Fields and Noble, *Int. J. Pept. Protein Res.* **35**:161-214 (1990)], or using automated synthesizers, such as sold by ABS. Thus, polypeptides of the invention may comprise D-amino acids, a combination of D- and L-amino acids, and various "designer" amino acids (e.g., β -methyl amino acids, α -methyl amino acids, and N^{α} -methyl amino acids, etc.) to convey special properties. Synthetic amino acids include ornithine for lysine, fluorophenylalanine for phenylalanine, and norleucine for leucine or isoleucine. Additionally, by assigning specific amino acids at specific coupling steps, α -helices, β turns, β sheets, γ -turns, and cyclic peptides can be generated.

Isolation and Crystallization of the Bacterial RNA Polymerase

The present invention provides a crystal of the Rif-RNAP complex that can be effectively diffract X-rays for the determination of the atomic coordinates of the Rif-RNAP to a resolution of better than 5.0 Angstroms and preferably to a resolution equal to or better than 3.5 Angstroms. The RNA polymerase can be expressed either

as described above or as described in U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entireties. Of course, the specific Rif-RNAP complex provided herein serves only as example, since the crystallization process can tolerate a broad range of active RNA polymerases and inhibitors. Therefore, any person with skill in the art of protein crystallization having the present teachings and without undue experimentation could crystallize a large number of alternative forms of the RNA polymerases from a variety of RNA polymerase fragments, or alternatively using a full length RNA polymerase from a related source and then allow the RNA polymerase to bind rifampicin and/or other

5 and inhibitors. Therefore, any person with skill in the art of protein crystallization having the present teachings and without undue experimentation could crystallize a large number of alternative forms of the RNA polymerases from a variety of RNA polymerase fragments, or alternatively using a full length RNA polymerase from a related source and then allow the RNA polymerase to bind rifampicin and/or other

10 RNAP binding partners (*e.g.*, inhibitors) as described below. As mentioned above, an RNA polymerase having conservative substitutions in its amino acid sequence are also included in the invention, including a selenomethionine substituted form.

Crystals of the RNA polymerase can be grown by a number of techniques including batch crystallization, vapor diffusion (either by sitting drop or hanging drop) and by

15 microdialysis. Seeding of the crystals in some instances is required to obtain X-ray quality crystals. Standard micro and/or macro seeding of crystals may therefore be used.

The crystals of the RNA polymerase can be grown alone or co-crystallized with a binding partner such as rifampicin. If the crystals are grown alone they can be

20 subsequently soaked in a stabilization buffer with an RNAP binding partner such as rifampicin (0.1 mM rifampicin was added in the Example below). The RNAP/RNAP-binding partner are preferably incubated in the stabilization buffer for at least twelve hours. An exemplary stabilization buffer contains between 1.7 - 2.3 M $(\text{NH}_4)_2\text{SO}_4$, 0.02-1 M Tris-HCl, pH 6.5-8.5, and approximately 20 mM MgCl_2 .

25 The crystals are then prepared for cryo-crystallography by soaking the RNAP/RNAP-binding partner complex in a stabilization buffer (*e.g.*, 2 M $(\text{NH}_4)_2\text{SO}_4$, 0.1 M Tris-HCl, pH 8.0, and 20 mM MgCl_2 containing 50% (w/v) sucrose) before flash freezing.

Aside from the methodology exemplified below, alternative methods may also be used to characterize the crystals. For example, crystals can be characterized by using X-rays produced in a conventional source (such as a sealed tube or a rotating anode) or using a synchrotron source. Methods of characterization include, but are not limited to, precision photography, oscillation photography and diffractometer data collection. Selenium-Methionine may be used, or alternatively a mercury derivative dataset (*e.g.*, using PCMB) could be used in place of the selenium-methionine derivatization.

Structural determinations can be performed by calculating Patterson maps using PHASES [Furey and Swaminathan, *Methods Enzymol.*, **277**:590-620 (1997)] for the ethyl-HgCl₂ and Ta₆Br₁₄ derivatives and using the Pb-derivative as native, for example. In the Example below, the native core RNAP structure [Zhang *et al.*, *Cell* **98**:811-824 (1999); U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entirety] was used as a starting model for rigid body refinement and positional refinement against the observed amplitudes from the Rif-RNAP complex crystal (F_o^{Rif}) using CNS [Adams *et al.*, *Proc. Natl. Acad. Sci. USA*, **94**:5018-5023 (1997)], yielding an initial R-factor of 0.354 ($R_{free} = 0.41$, where the same set of reflections was set aside as was used for the R_{free} determination of the native structure) for data from 100 - 3.2 Å resolution. An initial Fourier difference map, calculated using $|F_o^{Rif} - F_o^{nat}|$ amplitude coefficients and using phases calculated from the native core RNAP structure (ϕ^{nat}) clearly revealed density for the rifampicin molecule (Fig. 3a). Multiple rounds of manual rebuilding against $(2|F_o| - |F_c|)$ maps using O [Jones *et al.*, *Acta Cryst.*, **A 47**:110-119 (1991)], and refinement using CNS [Adams *et al.*, *Proc. Natl. Acad. Sci. USA*, **94**:5018-5023 (1997)] resulted in the current model (Table 1). At later stages of the refinement, the rifampicin X-ray crystal structure [Brufani *et al.*, *J. Molec. Biol.* **87**:409-435 (1974)] was placed into the difference density. Included in the model is the recently determined sequence of the *Taq* ω subunit modeled earlier as a polyalanine chain [Zhang *et al.*, *Cell* **98**:811-824 (1999); U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their

entireties]. Absent from the model is a 300 amino acid, non-conserved domain inserted between conserved regions A and B of the β' subunit [Zhang *et al.*, *Cell* **98**:811-824 (1999); U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entireties].

5 Protein-structure Based Design of Inhibitors of Bacterial RNA Polymerase

Once the three-dimensional structure of a crystal comprising a Rif-RNAP complex is determined, (*e.g.*, *see* the coordinates in Table 2 below, in Appendix following the Sequence Listing) a potential modulator of RNA Polymerase, can be examined through the use of computer modeling using a docking program such as GRAM,
 10 DOCK, or AUTODOCK [Dunbrack *et al.*, *Folding & Design*, **2**:27-42 (1997)], to identify potential modulators of the RNA Polymerase. This procedure can include computer fitting of potential modulators to the RNA Polymerase to ascertain how well the shape and the chemical structure of the potential modulator will bind to either the individual bound subunits or to the RNA Polymerase [Bugg *et al.*, *Scientific*
 15 *American*, **Dec.**:92-98 (1993); West *et al.*, *TIPS*, **16**:67-74 (1995)]. Computer programs can also be employed to estimate the attraction, repulsion, and steric hindrance of the subunits with a modulator/inhibitor (*e.g.*, the RNA Polymerase and a potential stabilizer).

Indeed, the shape of RNA polymerase resembles a crab-claw, with an internal groove
 20 or channel running along the full-length (between the claws). The molecule is about 150 Å long (from the back to the tips of the claws), 115 Å tall, and 110 Å wide (along the direction of the channel). The channel has many internal features, but the overall width is about 27 Å [*see*, U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entireties].

25 As disclosed herein the three-dimensional structure demonstrates that rifampicin binds the *Taq* core RNAP with a close complementary fit in a pocket between two structural domains of the RNAP β subunit. Only small, local conformational changes of both the inhibitor and the protein is observed. The binding site is deep within the

Importantly, the structural information disclosed herein demonstrates that rifampicin inhibits RNA polymerase by physically blocking transcription elongation. This is in direct contrast with the *modus operandi* of a classical enzyme inhibitor which generally binds to the catalytic center or with a key transition state intermediate. Therefore, the effect of rifampicin depends only on its ability to bind tightly to a relatively non-conserved part of the structure, thereby disrupting a critical RNAP function. Thus, the structural information disclosed herein provides the impetus to investigate the binding of other unrelated small molecules to any of a variety of sites within the RNAP channel, which could also block transcription elongation. A preferred site is one that is critical for the transcriptional activity of bacterial RNA polymerase, but one that is not required by the corresponding mammalian enzyme.

Towards this end, generally the tighter the fit, the lower the steric hindrances, and the greater the attractive forces, the more potent the potential modulator (*e.g.*, an inhibitor) since these properties are consistent with a tighter binding constant. Furthermore, the more specificity in the design of a potential drug the more likely that the drug will not interact as well with other proteins. This will minimize potential side-effects due to unwanted interactions with other proteins.

Initially alternative compounds known to bind bacterial RNA polymerase, including rifampicin analogs, can be systematically modified by computer modeling programs until one or more promising potential analogs are identified. In addition systematic modification of selected analogs can then be systematically modified by computer modeling programs until one or more potential analogs are identified. Such analysis has been shown to be effective in the development of HIV protease inhibitors [Lam *et al.*, *Science* **263**:380-384 (1994); Wlodawer *et al.*, *Ann. Rev. Biochem.* **62**:543-585 (1993); Appelt, *Perspectives in Drug Discovery and Design* **1**:23-48 (1993); Erickson, *Perspectives in Drug Discovery and Design* **1**:109-128 (1993)]. Alternatively a

5

10

20

25

embodiment, isothermal calorimetry can be used to determine the stability of the bacterial RNA polymerase in the absence and presence of the potential modulator.

In another embodiment, a Biacore machine can be used to determine the binding constant of the bacterial RNA polymerase to a DNA template in the presence and
 5 absence of the potential modulator. Alternatively, one or more of the bacterial RNA polymerase subunits can be immobilized on a sensor chip. The remaining subunits can then be contacted with (*e.g.*, flowed over) the sensor chip to form the bacterial RNA polymerase.

In this case the dissociation constant for the bacterial RNA polymerase can be
 10 determined by monitoring changes in the refractive index with respect to time as buffer is passed over the chip [O'Shannessy *et al.* *Anal. Biochem.* **212**:457-468 (1993); Schuster *et al.*, *Nature* **365**:343-347 (1993)]. Scatchard plots, for example, can be used in the analysis of the response functions using different concentrations of a particular subunit. Flowing a potential modulator at various concentrations over the
 15 bacterial RNA polymerase and monitoring the response function (*e.g.*, the change in the refractive index with respect to time) allows the bacterial RNA polymerase dissociation constant to be determined in the presence of the potential modulator and thereby indicates whether the potential modulator is either an inhibitor, or an agonist of the bacterial RNA polymerase complex.

20 In another aspect of the present invention a potential modulator is assayed for its ability to inhibit the bacterial RNA polymerase. A modulator that inhibits the RNA polymerase can then be selected. In a particular embodiment, the effect of a potential modulator on the catalytic and/or transcriptional activity of bacterial RNA polymerase is determined. The potential modulator is then be added to a bacterial culture to
 25 ascertain its effect on bacterial proliferation. A potential modulator that inhibits bacterial proliferation can then be selected.

In a particular embodiment, the effect of the potential modulator on the catalytic and/or transcriptional activity of the bacterial RNA polymerase is determined (either independently, or subsequent to a binding assay as exemplified above). In one such embodiment, the rate of the DNA-dependent RNA transcription is determined. For such assays a labeled nucleotide could be used. This assay can be performed using a real-time assay *e.g.*, with a fluorescent analog of a nucleotide. Alternatively, the determination can include the withdrawal of aliquots from the incubation mixture at defined intervals and subsequent placing of the aliquots on nitrocellulose paper or on gels. In a particular embodiment the potential modulator is selected when it is an inhibitor of the bacterial RNA polymerase.

One assay for RNA polymerase activity is a modification of the method of Burgess *et al.* [*J. Biol. Chem.*, **244**:6160 (1969)]

[See also <http://www.worthington-biochem.com/manual/R/RNAP.html>].

One unit incorporates one nanomole of UMP into acid insoluble products in 10 minutes at 37°C under the assay conditions such as those listed below.

The suggested reagents are:

- (a) 0.04 M Tris-HCl, pH 7.9, containing 0.01 M MgCl₂, 0.15 M KCl, and 0.5 mg/ml BSA;
- (b) Nucleoside triphosphates (NTP) : 0.15 mM each of ATP, CTP, GTP, UTP; spiked with ³H - UTP 75000 - 150000 cpms/0.1 ml;
- (c) 0.15 mg/ml calf thymus DNA;
- (d) 10% cold perchloric acid; and
- (e) 1% cold perchloric acid.

0.1 - 0.5 units of RNA polymerase in 5 µl - 10 µl is used as the starting enzyme concentration.

The procedure is to add 0.1 ml Tris-HCl, 0.1 ml NTP and 0.1 ml DNA to a test tube for each sample or blank. At zero time enzyme (or buffer for blank) is added to each

test tube, and the contents are then mixed and incubated at 37°C for 10 minutes. 1 ml of 10% perchloric acid is added to the tubes to stop the reaction. The acid insoluble products can be collected by vacuum filtration through MILLIPORE filter discs having a pore size of 0.45 u - 10 u (or equivalent). The filters are then washed four
 5 times with 1% cold perchloric acid using 1 ml - 3 ml for each wash. These filters are then placed in scintillation vials. 2 mls of methyl cellosolve are added to the scintillation vials to dissolve the filters. When the filters are completely dissolved (after about five minutes) 10 mls of scintillation fluid are added and the vials are counted in a scintillation counter.

- 10 For calculation of units of RNA polymerase/mg of protein the following equation can be used:

$$15 \quad \text{units/mg} = \frac{\text{CPM}_{\text{test}} - \text{CPM}_{\text{blank}}}{\text{CPM}_{\text{total}} \times \text{mg protein}_{\text{in test}}}$$

Alternative transcription assays can also be employed [*see* Example below, and Nudler *et al.*, *Science* **265**:793-796 (1994)]. One such assay comprises a core RNAP that can be incubated with a suitable σ subunit to form the holoenzyme. A potential modulator can then be added prior to, simultaneously with, subsequently to a
 20 promoter fragment (*e.g.*, T7A1 as exemplified below). RNA synthesis is then initiated by the addition of a primer (*e.g.*, a CpA primer) and the four nucleotide triphosphates (NTPs). The RNA synthesis in the presence and absence of the potential modulator is then quantified. In the Example below, a radioactive nucleotide was employed and the radioactive RNA products were analyzed on a 15%
 25 polyacrylamide sequencing gel. Alternatively, a fluorescent nucleotide analog can be used. Transcription reactions on a minimal scaffold system can be performed as shown in Fig. 6b below in the presence and the absence of the potential modulator [*see also* Korzheva *et al.*, *Science* **289**:619-625 (2000)].

When suitable potential modulators are identified, a supplemental crystal can be
 30 prepared which comprises the bacterial RNA polymerase and the potential modulator

(see Example below). Preferably the crystal effectively diffracts X-rays for the determination of the atomic coordinates of the protein-ligand complex to a resolution of better than 5.0 Angstroms, more preferably equal to or better than 3.5 Angstroms.

The three-dimensional structure of the supplemental crystal can be determined by

- 5 Molecular Replacement Analysis. Molecular replacement involves using a known three-dimensional structure as a search model to determine the structure of a closely related molecule or protein-ligand complex in a new crystal form. The measured X-ray diffraction properties of the new crystal are compared with the search model structure to compute the position and orientation of the protein in the new crystal.
- 10 Computer programs that can be used include: X-PLOR (see above), CNS, (Crystallography and NMR System, a next level of XPLOR), and AMORE [J. Navaza, *Acta Crystallographica ASO*, 157-163 (1994)]. Once the position and orientation are known an electron density map can be calculated using the search model to provide X-ray phases. Thereafter, the electron density is inspected for
- 15 structural differences and the search model is modified to conform to the new structure. Using this approach, it is also possible to use the claimed crystal of the Rif-RNAP complex to solve the three-dimensional structures of other bacterial core RNA polymerases bound to rifampicin (and/or other inhibitors) having pre-ascertained amino acid sequences. Other computer programs that can be used to solve the
- 20 structures of the bacterial RNA polymerase from other organisms include: QUANTA, CHARMM; INSIGHT; SYBYL; MACROMODE; and ICM.

A candidate drug can be selected by performing rational drug design with the three-dimensional structure determined for the supplemental crystal, preferably in conjunction with computer modeling discussed above. The candidate drug (*e.g.*, a

25 potential modulator of bacterial RNA polymerase) can then be assayed as exemplified above, or *in situ*. A candidate drug can be identified as a drug, for example, if it inhibits bacterial proliferation.

A potential inhibitor (*e.g.*, a candidate drug) would be expected to interfere with bacterial growth. Therefore, an assay that can measure bacterial growth may be used to identify a candidate drug.

Methods of testing a potential bactericidal agent (*e.g.*, the candidate drug) in an animal model are well known in the art, and can include standard bactericidal assays. The potential modulators can be administered by a variety of ways including topically, orally, subcutaneously, or intraperitoneally depending on the proposed use. Generally, at least two groups of animals are used in the assay, with at least one group being a control group which is administered the administration vehicle without the potential modulator.

For all of the drug screening assays described herein further refinements to the structure of the drug will generally be necessary and can be made by the successive iterations of any and/or all of the steps provided by the particular drug screening assay.

Labels

Suitable labels include enzymes, fluorophores *e.g.*, fluorescein isothiocyanate (FITC), phycoerythrin (PE), Texas red (TR), rhodamine, free or chelated lanthanide series salts, especially Eu^{3+} , to name a few fluorophores and including fluorescent GTP and GDP analogs such as mantGTP and mantGDP, chromophores, radioisotopes, chelating agents, dyes, colloidal gold, latex particles, ligands (*e.g.*, biotin), and chemiluminescent agents. When a control marker is employed, the same or different labels may be used for the test and control marker.

In the instance where a radioactive label, such as the isotopes ^3H , ^{14}C , ^{32}P , ^{35}S , ^{36}Cl , ^{51}Cr , ^{57}Co , ^{58}Co , ^{59}Fe , ^{90}Y , ^{125}I , ^{131}I , and ^{186}Re are used, known currently available counting procedures may be utilized. In the instance where the label is an enzyme, detection may be accomplished by any of the presently utilized colorimetric,

spectrophotometric, fluorospectrophotometric, amperometric or gasometric techniques known in the art.

Direct labels are one example of labels which can be used according to the present invention. A direct label has been defined as an entity, which in its natural state, is readily visible, either to the naked eye, or with the aid of an optical filter and/or applied stimulation, *e.g.* ultraviolet light to promote fluorescence. Among examples of colored labels, which can be used according to the present invention, include metallic sol particles, for example, gold sol particles such as those described by Leuving (U.S. Patent 4,313,734); dye sol particles such as described by Gribnau *et al.* (U.S. Patent 4,373,932) and May *et al.* (WO 88/08534); dyed latex such as described by May, *supra*, Snyder (EP-A 0 280 559 and 0 281 327); or dyes encapsulated in liposomes as described by Campbell *et al.* (U.S. Patent 4,703,017) Other direct labels include a radionucleotide, a luminescent moiety, or a fluorescent moiety including as a modified/fusion chimera of green fluorescent protein (as described in U.S. Patent No. 5,625,048 filed April 29, 1997, and WO 97/26333, published July 24, 1997, the disclosures of each are hereby incorporated by reference herein in their entireties). In addition to these direct labeling devices, indirect labels comprising enzymes can also be used according to the present invention. Various types of enzyme linked immunoassays are well known in the art, for example, alkaline phosphatase and horseradish peroxidase, lysozyme, glucose-6-phosphate dehydrogenase, lactate dehydrogenase, urease, these and others have been discussed in detail by Eva Engvall in Enzyme Immunoassay ELISA and EMIT in *Methods in Enzymology*, **70**:419-439 (1980) and in U.S. Patent 4,857,453.

Suitable enzymes include, but are not limited to, alkaline phosphatase and horseradish peroxidase. Other labels for use in the invention include magnetic beads or magnetic resonance imaging labels.

In another embodiment, a phosphorylation site can be created on an antibody of the invention for labeling with ^{32}P , *e.g.*, as described in European Patent No. 0372707

(application No. 89311108.8) by Sidney Pestka, or U.S. Patent No. 5,459,240, issued October 17, 1995 to Foxwell *et al.*

As exemplified herein, proteins, including antibodies, can be labeled by metabolic labeling. Metabolic labeling occurs during *in vitro* incubation of the cells that express the protein in the presence of culture medium supplemented with a metabolic label, such as [³⁵S]-methionine or [³²P]-orthophosphate. In addition to metabolic (or biosynthetic) labeling with [³⁵S]-methionine, the invention further contemplates labeling with [¹⁴C]-amino acids and [³H]-amino acids (with the tritium substituted at non-labile positions).

Three-Dimensional Representation of the Structure of the Rif-RNAP complex

In addition, the present invention provides a computer that comprises a representation of the RNAP-RNAP binding partner complex (*e.g.*, the Rif-RNAP complex) in computer memory that can be used to screen for compounds that will or are likely to inhibit RNAP. In a related embodiment, the computer can be used in the design of altered RNAPs that have either enhanced, or alternatively diminished RNA polymerase activity. Preferably, the computer comprises portions of and/or all of the information contained in Table 2. In a particular embodiment, the computer comprises: (i) a machine-readable data storage material encoded with machine-readable data, (ii) a working memory for storing instructions for processing the machine readable data, (iii) a central processing unit coupled to the working memory and the machine-readable data storage material for processing the machine-readable data into a three-dimensional representation, and (iv) a display coupled to the central processing unit for displaying the three-dimensional representation.

Thus the machine-readable data storage medium comprises a data storage material encoded with machine readable data which can comprise portions and/or all of the structural information contained in Table 2. One embodiment for manipulating and displaying the structural data provided by the present invention is schematically depicted in Figure 7. As depicted, the System 1, includes a computer 2 comprising a

central processing unit ("CPU") 3, a working memory 4 which may be random-access memory or "core" memory, mass storage memory 5 (e.g., one or more disk or CD-ROM drives), a display terminal 6 (e.g., a cathode-ray tube), one or more keyboards 7, one or more input lines 10, and one or more output lines 20, all of which are
5 interconnected by a conventional bidirectional system bus 30.

Input hardware 12, coupled to the computer 2 by input lines 10, may be implemented in a variety of ways. Machine-readable data may be inputted *via* the use of one or more modems 14 connected by a telephone line or dedicated data line 16.

Alternatively or additionally, the input hardware 12 may comprise CD-ROM or disk
10 drives 5. In conjunction with the display terminal 6, the keyboard 7 may also be used as an input device. Output hardware 22, coupled to computer 2 by output lines 20, may similarly be implemented by conventional devices. Output hardware 22 may include a display terminal 6 for displaying the three dimensional data. Output hardware might also include a printer 24, so that a hard copy output may be produced,
15 or a disk drive 5, to store system output for later use, *see also* U.S. Patent No: 5,978,740, Issued November 2, 1999, the contents of which are hereby incorporated by reference in their entirety.

In operation, the CPU 3 (i) coordinates the use of the various input and output devices 12 and 22; (ii) coordinates data accesses from mass storage 5 and accesses to and
20 from working memory 4; and (iii) determines the sequence of data processing steps. Any of a number of programs may be used to process the machine-readable data of this invention.

The present invention may be better understood by reference to the following non-limiting Example, which is provided as exemplary of the invention. The following
25 example is presented in order to more fully illustrate the preferred embodiments of the invention. It should in no way be construed, however, as limiting the broad scope of the invention.

EXAMPLE
STRUCTURAL MECHANISM FOR RIFAMPICIN INHIBITION
OF BACTERIAL RNA POLYMERASE

Introduction

5 High-resolution structural studies of the Rif-RNAP complex should lead to insights into rifampicin binding, the mechanism of inhibition, and also the mechanism by which mutations lead to Rif^R. These structural studies will also shed light on the transcription mechanism itself, as well as provide the basis for the development of drugs that selectively inhibit bacterial RNAPs, but are less prone than rifampicin to
10 lead bacterial mutations/substitutions of single amino acids that give rise to resistance. Indeed, the recent determination of the crystal structure of core RNAP from *Thermus aquaticus* (Taq) [Zhang *et al.*, *Cell* **98**:811-824 (1999); U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entirety] has opened the door to further studies of RNAP structure, function,
15 and interactions with substrates, ligands, and inhibitors.

To further provide a more detailed framework to interpret the existing genetic, biochemical, and biophysical information, as well as to guide further studies aimed at understanding the transcription process and its regulation, the three-dimensional structure of a bacterial core RNAP complexed with rifampicin was determined by
20 X-ray crystallography at 3.3 Å resolution as detailed below. The structure explains the effects of rifampicin on RNAP function. In combination with a model of the ternary transcription complex and biochemical experiments, the data indicate that the predominant effect of rifampicin on RNAP function is to directly block the path of the elongating RNA transcript at the 5'-end when the transcript becomes either 2 or 3
25 nucleotides in length.

Methods

Purification and crystallization: Native Taq core RNAP was purified and crystallized as described previously [Zhang *et al.*, *Cell* **98**:811-824 (1999); U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby

- incorporated by reference in their entireties]. Crystals were subsequently soaked in stabilization solution [2 M $(\text{NH}_4)_2\text{SO}_4$, 0.1 M Tris-HCl, pH 8.0, and 20 mM MgCl_2] with 0.1 mM rifampicin for at least 12 hours. The crystals were then prepared for cryo-crystallography by soaking in stabilization solution containing 50% (w/v) sucrose for 30 minutes before flash freezing in liquid nitrogen. Diffraction data was collected at the APS beamline SBC 19ID using 0.3° oscillations, and processed using DENZO and SCALEPACK [Otwinowski, *Isomorphous Replacement and Anomalous Scattering* (eds. Wolf, Evans and Leslie) Science and Engineering Research Council, Daresbury Laboratory, Daresbury, UK, (1991)].
- 10 In short, the preparative procedure for *T. aquaticus* core RNAP is similar to the preparation of *E. coli* core RNAP [Polyakov *et al.*, *Cell*, **83**:365-373 (1995)]. Briefly, approximately 200 g wet cell paste is thawed and lysed using a continuous-flow French press. After a low-speed spin, the soluble fraction is precipitated with 0.6% Polymin-P. RNAP is eluted from the Polymin-P pellet with TGED buffer (10 mM
- 15 Tris -HCl, pH 8, 5% glycerol, 1 mM EDTA, 1 mM DTT) plus 1 M NaCl, then precipitated by adding 33%(g/v) ammonium sulfate. The pellet is resuspended and loaded onto a 50 ml column of heparin-SEPHAROSE FF (Pharmacia) equilibrated with TGED buffer *plus* 0.2 M NaCl. The RNAP is eluted from the column with TGED buffer *plus* 0.6 M NaCl. The RNAP was again precipitated with ammonium
- 20 sulfate, then resuspended and loaded on a SUPERDEX-200 gel filtration column equilibrated with TGED buffer *plus* 0.5 M NaCl. Fractions containing RNAP were pooled and loaded onto a MONO-Q (Pharmacia) ion-exchange column equilibrated with TGED buffer *plus* 0.1 M NaCl. The protein was eluted with a gradient from 0.1 to 0.5 M NaCl. The RNAP peak eluted at around 0.3 M NaCl. The RNAP was
- 25 concentrated using a centrifugal filter, then loaded onto an SP SEPHAROSE (Pharmacia) column equilibrated in TGED buffer *plus* 0.1 M NaCl. After loading, the column was incubated at 4°C for at least 10 hours, then pure RNAP was eluted with a 0.1 to 0.5 M NaCl gradient (core RNAP elutes at around 0.3 M NaCl). 200 g wet cell paste typically yielded 15 mg of core RNAP, which was more than 99% pure as

judged from overloaded, Coomassie-stained SDS gels. This sample is ready for crystallization.

Crystals of *T. aquaticus* core RNAP were grown by vapor diffusion. 10 μ l of *T. aquaticus* core RNAP (17 mg/ml) was mixed with the same volume of a solution
 5 containing 40-45% saturated $(\text{NH}_4)_2\text{SO}_4$, 0.1 M Tris-HCl, pH 8.0, and 20 mM MgCl_2 , and incubated as a hanging drop over the same solution. Crystals grow in 2-3 weeks to typical dimensions of 0.15 mm X 0.15 mm X 0.4 mm at room temperature. For cryo-crystallography, the crystals are pre-soaked in stabilization solution (same as the crystallization solution except with 50% saturated ammonium sulfate). The crystals
 10 are then soaked in stabilization solution containing 50% (g/v) sucrose for about 30 minutes before flash freezing. The frozen crystals diffract to 5.0 Å from an in-house X-ray generator. Spots can sometimes be observed, in one direction, to 2.7 Å resolution at synchrotron beamlines. Diffraction data was processed using DENZO and SCALEPACK [Otwinowski, *Isomorphous Replacement and Anomalous*
 15 *Scattering* (eds. Wolf, Evans and Leslie) Science and Engineering Research Council, Daresbury Laboratory, Daresbury, UK, (1991)].

Selenomethionyl core RNAP was prepared and crystallized using the same procedures from *T. aquaticus* cells grown in minimal media (culture medium 162) [Degryse *et al.*, *Arch. Microbiol.*, **117**:189-196 (1978)]. Cells were induced to incorporate
 20 selenomethionine by suppression of methionine biosynthesis [Doublie, *Methods Enzymol.*, **276**:523-530 (1997)].

Structure Determination: The native core RNAP structure [Zhang *et al.*, *Cell* **98**:811-824 (1999); U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entirety] was used as
 25 a starting model for rigid body refinement and positional refinement against the observed amplitudes from the Rif-RNAP complex crystal (F_o^{Rif}) using CNS [Adams *et al.*, *Proc. Natl. Acad. Sci. USA*, **94**:5018-5023 (1997)], yielding an initial R-factor of 0.354 ($R_{\text{free}} = 0.41$, where the same set of reflections was set aside as was used for

the R_{free} determination of the native structure) for data from 100 - 3.2 Å resolution. An initial Fourier difference map, calculated using $|F_o^{\text{Rif}} - F_o^{\text{nat}}|$ amplitude coefficients and using phases calculated from the native core RNAP structure (ϕ^{nat}) clearly revealed density for the Rif molecule (Fig. 3a). Multiple rounds of manual rebuilding against
 5 $(2|F_o| - |F_c|)$ maps using O [Jones *et al.*, *Acta Cryst.*, **A 47**:110-119 (1991)], and refinement using CNS [Adams *et al.*, *Proc. Natl. Acad. Sci. USA*, **94**:5018-5023 (1997)] resulted in the current model (Table 1). At later stages of the refinement, the Rif X-ray crystal structure [Brufani *et al.*, *J. Molec. Biol.* **87**:409-435 (1974)] was easily placed into the difference density. Included in the model is the recently
 10 determined sequence of the *Taq* ω subunit modeled earlier as a polyalanine chain [Zhang *et al.*, *Cell* **98**:811-824 (1999); U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entirety]. Absent from the model is a 300 amino acid, non-conserved domain inserted between conserved regions A and B of the β' subunit [Zhang *et al.*, *Cell*
 15 **98**:811-824 (1999); U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entirety].

Assays: *Taq* cells were tested for sensitivity to rifampicin on solid media. Plates containing 3% bactoagar and 1/5 dilution of Luria broth were poured with and without 50 µg/ml of rifampicin. Cells from frozen stock were then streaked onto plates and
 20 incubated at 65°C for 2 days and assessed for growth.

The transcription assay comparing rifampicin inhibition of *E.coli* and *Taq* RNAPs (Fig. 2a) was performed as previously described [Nudler *et al.*, *Science* **265**:793-796 (1994)]. Briefly, 0.1 pmol of purified *Taq* core RNAP [Zhang *et al.*, *Cell* **98**:811-824 (1999); U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which
 25 are hereby incorporated by reference in their entirety] was incubated with *Taq* σ^A in 20 µl of transcription buffer (40 mM Tris-HCl, pH 7.9, 40 mM KCl, 5 mM MgCl₂) for 15 minutes at 37°C to form holoenzyme. Rifampicin was added to the final concentrations indicated in Fig. 2a and incubated another 5 minutes at 37°C, followed by the addition of 0.15 pmol of T7A1 promoter fragment and incubation for 5 minutes

at 37°C. RNA synthesis was initiated by the addition of CpA primer (100 µM), NTPs (25 µM each), and α-[³²P]UTP (0.3 µM), and the reaction was stopped after incubation for 10 minutes at 37°C. The assay for *E.coli* RNAP holoenzyme was the same except the CpA primer was added to a concentration of 10 µM. Radioactive RNA products were analyzed on a 15% polyacrylamide sequencing gel.

Assays for extension of the Rif-nucleotide compounds (Fig. 2c-2d) were carried out as described [Mustaev *et al.*, *Proc.Nat.Acad.Sci.USA* **91**:12036-12040 (1994)] with minor modifications. After binary complex formation, transcription reactions were started by the addition 10 µM Rif-(CH₂)_n-A compound, with the 'n' indicated in Fig. 2c-2d, and α-[³²P]UTP (0.3 µM). The reactions were incubated for 2 minutes at room temperature for *E.coli* RNAP and 3 minutes at 55°C for *Taq*. Under these conditions, the reaction was not complete, and the yield of the Rif-(CH₂)_n-ApU depended on the linker length. Radioactive RNA products were analyzed on a 23% polyacrylamide sequencing gel.

Transcription reactions on the minimal scaffold system shown (Fig. 6b) were performed as described [Korzheva *et al.*, *Science* **289**:619-625 (2000)] with minor modifications. The RNA and DNA components of the scaffold (100 pmol of each) were mixed in 100 µl of transcription buffer at 45°C and the mixture was allowed to cool to room temperature over 30 minutes. RNAP/scaffold complexes were formed by incubation of the annealed scaffold (10 pmol) with a molar equivalent of core RNAP (either *E.coli* or *Taq*) which was preincubated with rifampicin (100 µM for *E.coli*, 200 µM for *Taq*) for 10 minutes to form the RNAP/scaffold complex. Extension of the RNA oligonucleotide was assayed by the addition of α-[³²P]CTP (0.3 µM) and a 5 minute incubation at room temperature. In Fig. 6b, lanes 1-5 and 16-20, RNAP was preincubated with rifampicin (100 µM for *E. coli* RNAP, 200 µM for *Taq*) for 10 minutes. In lanes 6-10 and 21-25, the RNAP/scaffold complexes formed in the absence of rifampicin were incubated with rifampicin (concentrations as above) for 10 minutes. Finally, in lanes 11-15 and 26-30, the RNAP or RNAP/scaffold complex

was not exposed to rifampicin. Radioactive RNA products were analyzed on a 23% polyacrylamide sequencing gel.

Results

Rifampicin Inhibition of Taq RNAP: From a biochemical perspective, the interaction of rifampicin (Rif) with RNAP has been extensively characterized using *E. coli* RNAP, which served as a prototype for bacterial pathogens [Drancourt and Raoult, *Antimicrob. Agents Chemother.* **43**:2400-2403 (1999); Heep *et al.*, *Antimicrob. Agents Chemother.* **44**:1075-1077 (1999); Honore *et al.*, *Molec. Microbiol.* **7**:207-214 (1993); Morse *et al.*, *J. Clin. Microbiol.* **37**:2913-2929 (1999); Nolte, *J. Antimicrob. Chemother.* **39**:747-755 (1997); Padayachee and Klugman, *Antimicrob. Agents Chemother.* **43**:2361-2365 (1999); Ramaswamy and Musser, *Tubercle and Lung Disease* **79**:3-29 (1998); Wichelhaus *et al.*, *Antimicrob. Agents Chemother.* **43**:2813-2816 (1999)]. The inhibition of *Taq* RNAP by rifampicin was therefore investigated to assess this system as a structural model for Rif-RNAP interactions.

Sequence comparisons in the four distinct regions of *rpoB* which harbor Rif^R mutations indicate a very high level of conservation among prokaryotes. Between *E. coli*, *Taq*, and *M. tuberculosis*, the sequences are 91% identical over 60 residues (93% conserved), explaining the broad spectrum of rifampicin activity. Nevertheless, among the 23 positions with single amino-acid substitutions that give rise to Rif^R in either *E. coli* or *M. tuberculosis*, 5 of these positions (*Taq* β 387, 395, 398, 453, and 566; the *Taq* numbering is used throughout this application unless otherwise specified) are substituted in *Taq* (Fig. 1). In contrast, there is a relatively low level of conservation between prokaryotes and eukaryotes within these regions (Fig. 1), explaining the lack of rifampicin activity against eukaryotic RNAPs and eukaryotic cells.

A plate assay (*see* Methods above) was used to show that *Taq* cells were unable to grow on media supplemented with 50 μ g/ml rifampicin. For *in vitro* studies, *Taq* RNAP holoenzyme was reconstituted using *Taq* core RNAP purified from *Taq* cells [Zhang *et al.*, *Cell* **98**:811-824 (1999); U.S. Serial No.09/396,651, Filed September

15, 1999, the contents of which are hereby incorporated by reference in their entireties] and recombinant *Taq* σ^A (overexpressed and purified from *E. coli*). The enzyme initiated, elongated, and terminated transcripts efficiently from a template containing the T7A1 promoter and the tR2 intrinsic terminator (Fig. 2a) [Nudler *et al.*, 5 *J.Molec.Biol.* **288**:1-12(1994)] at 37°C using the dinucleotide CpA as the initiating primer. The major RNA products, a trimeric abortive transcript (CpApU), a 105 nucleotide terminated transcript (Term), and a 127 nucleotide runoff transcript (Run off), were the same as those produced by *E. coli* RNAP (Fig. 2a, lanes 1 and 8). Since *E. coli* σ^{70} is totally inactive when combined with *Taq* core RNAP in this assay, the possibility of trace contamination with *E. coli* σ^{70} does not affect the conclusions from this assay for *Taq* RNAP. Quantitatively, the two RNAPs responded very differently to rifampicin, the K_i (estimated from the rifampicin concentration where the production of long transcripts was inhibited by 50%) for *E. coli* RNAP was about 0.1 μ M, while for *Taq* RNAP it was about 10 μ M, a 100-fold difference in sensitivity. 10 Qualitatively, however, both RNAPs responded the same way, with an increase in the production of the trimeric product and a concurrent precipitous drop in the production of the long transcripts (Fig. 2a). 15

Mustaev *et al.*, [*Proc.Nat.Acad.Sci.USA* **91**:12036-12040 (1994)] used chimeric Rif-nucleotide compounds to measure the distance between the initiating nucleotide binding site (the i-site) and the Rif binding site. By varying the linker between the Rif 20 and the nucleotide and testing for maximal transcription initiation activity, the optimal length was found that allowed binding of each moiety in its respective site. This experiment was used to compare the disposition of the Rif and i-sites in *E. coli* and *Taq* RNAP. In both cases, optimal initiation activity was observed when the linker 25 comprised five -(CH₂)- groups (Figs. 2c-2d). Thus, in spite of the fact that *Taq* RNAP requires a 100-fold higher concentration of rifampicin for inhibition, *Taq* RNAP binds rifampicin and is inhibited through the same biochemical mechanism as *E. coli* RNAP, and the disposition of the Rif-site with respect to the universally conserved active site is identical. Therefore, *Taq* RNAP can serve as a model for 30 rifampicin interactions with other RNAPs.

Rif-RNAP Structure Determination and Refinement: Tetragonal crystals of *Taq* core RNAP [Zhang *et al.*, *Cell* **98**:811-824 (1999); U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entireties] were incubated overnight in stabilization buffer with 0.1 mM rifampicin, followed by a 30 minute soak in cryo-solution (without rifampicin) before flash freezing. During this procedure, the crystals took on a deep orange color, confirming the binding of rifampicin. The same results were obtained with co-crystals grown in the presence of 0.1 mM rifampicin, suggesting that rifampicin binding causes few if any conformational changes in the RNAP.

- 10 The *Taq* core RNAP:Rif crystals were isomorphous with the native *Taq* core RNAP crystals [Zhang *et al.*, *Cell* **98**:811-824 (1999); U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entireties]. Strong electron density was observed in difference Fourier maps for the rifampicin (Fig. 3a), which occupies a shallow pocket between β structural domains 3 and 4 (Fig. 3b) that is surrounded by the known Rif^R mutations (Fig. 1)
- 15 [Zhang *et al.*, *Cell* **98**:811-824 (1999); U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entireties]. The electron difference density also indicated shifts and/or ordering of several β residues interacting directly with rifampicin, including Q390, L391, Q393, D396, H406, R409, and L413 (Fig. 4). Only very small shifts in localized regions of
- 20 the protein backbone were indicated.

The rifampicin X-ray crystal structure [Brufani *et al.*, *J. Molec.Biol.* **87**:409-435 (1974)] was easily placed into the difference density. Subsequent refinements resulted in only small shifts of the ansa chain (Fig. 3c) to better fit the density.

- 25 Multiple rounds of manual rebuilding against $(2|F_o| - |F_c|)$ maps and refinement resulted in the current model (*see* Methods above and Table 1).

Table 1

CRYSTALLOGRAPHIC DATA AND STRUCTURAL MODEL

<u>DIFFRACTION DATA</u>					
5	<u>Parameter</u>		<u>Total</u>		<u>Outer Shell</u>
	Resolution range (Å)		30-3.3		3.42-3.3
	Rmerge1 (%)		7.7		34.4
	Completeness (%)		86.1		71.7
	I/σI		10.7		1.7
10	No. of reflections		75,420		6,173
	No. of unique obs.		214,453		11,549
<hr/>					
<u>STRUCTURAL MODEL</u>					
<u>Number of Residues</u>					
15	Protein	Mr (kDa)	sequence	model	regions modeled
	Subunit ²				
	β'	170.7	1,525	1,139	3-31, 69-155 (poly-Ala), 452-523, 536-1241, 1250-1410, 1414-1497
20	β	124.4	1,119	1,114	2-1115
	αI	34.9	313	223	6-228
	αII	34.9	313	229	3-231
	ω	11.6	99	98	1-98
25	Total	376.5	3,369	2,803	
<hr/>					
REFINEMENT					
	R _{cryst} (%)	28.1			
30	R _{free} (%)	35.9			

$$^1\text{Rmerge} = \Sigma |I_j - \langle I \rangle| / \Sigma I_j$$

²Also included in the model was one Mg²⁺ and one Zn²⁺ ion [Zhang *et al.*, *Cell* 98:811-824 (1999); U.S. Serial No. 09/396,651, Filed September 15, 1999] and one

35 Rif molecule [Brufani *et al.*, *J. Molec. Biol.* 87:409-435 (1974)].

carboxyl of F394, respectively. O8 of rifampicin is also positioned to make a potential hydrogen bond with the backbone amide of F394.

- D396 contributes to the binding interface in several ways. In addition to forming a potential hydrogen bond with O10 of rifampicin, it forms the top end of the binding pocket (in Figs. 4a-4b) by making van-der-Waals contact with C18-C21, and C31. Moreover, the negative charge of D396 may be important for neutralizing the positive charges of two nearby side chains, R405 and R409 (Figs. 4a-4b), each about 6 Å away. The charge neutralization might be important for the binding of the relatively apolar of rifampicin. Most Rif^R mutants at amino acid residue 396 substitute a large, bulky group that would likely interfere with rifampicin binding and would not have the correct geometry for hydrogen bonding O10 (Y), or else substitute an apolar group (V, G, or A) with no hydrogen bonding ability. One of these mutants, D396V (amino acid position 516 in *E. coli*), was among the original, strong Rif^R mutants mapped by Ovchinnikov *et al.* [*Molec. Gen. Genet.* **190**:344-348 (1983)], pointing to the importance of this residue in forming the rifampicin binding interface. Another mutant identified in *E. coli*, however (D396N), is isosteric with aspartic acid and would likely maintain the hydrogen bond with O10. Nevertheless, this substitution yields weak Rif^R [Lisitsyn *et al.*, *Bioorg Khim* **10**:127-128 (1984)], which is likely caused by the loss of the negative charge at this position.
- Rifampicin has a partial +-charge, localized at N4 (Fig. 3c). A negatively-charged residue, E445, is situated nearby and may contribute to the rifampicin binding site by neutralizing this charge. This is not likely to be a strong effect, as many rifampicin derivatives with equal or stronger activity than rifampicin do not have this partial charge. E445 is the only residue close enough to rifampicin to be involved in potentially direct interactions (Figs. 4a-4b) for which a Rif^R mutant has not been reported. However, this residue is universally conserved as either glutamic acid or aspartic acid in a segment of β region D that is invariantly present in prokaryotes, chloroplast, archaeobacteria, and eukaryotes [Allison *et al.*, *Cell* **42**:599-610 (1985);

Sweetser *et al.*, *Proc.Natl.Acad.Sci.USA* **84**:1192-1196 (1987)], pointing to its importance for the basic function of RNAP.

Thus, of the 12 residues that are close enough to rifampicin to make direct interactions (including backbone interactions with F394; Figs. 4a-4b), 11 mutate to a Rif^R phenotype. The twelfth position, E445, is highly conserved so that its substitution would likely be lethal and consequently not be detectable as a Rif^R mutation.

Twelve additional positions have been identified at which substitution gives rise to Rif^R (Fig. 1). These residues surround the Rif binding pocket but do not make direct interactions with the antibiotic (Figs. 5a-5b). In every case, the Rif^R mutations involve replacement by a different sized amino acid side-chain (almost always substituting a small residue with a more bulky one), or else involve adding or removing a proline residue. These substitutions would likely affect the folding or packing of the protein in the local vicinity of the substituted residue, causing distortions of the Rif binding pocket.

Mechanism of RNAP Inhibition by Rif: The effects of rifampicin on RNAP in each stage of the transcription cycle have been probed using detailed kinetic analyses. Rifampicin has essentially no effect on specific promoter binding and open complex formation [Hinkle *et al.*, *J.Molec.Biol.*, **70**, 209-220 (1972); McClure and Cech, *J.Biol.Chem.* **253**:8949-8956 (1978)]. A small increase (about 2-fold) in the apparent *K_m* for initiating substrate binding in the enzyme's i-site (the 5'-nucleotide) was observed, but the binding of the incoming nucleotide substrate in the i+1 site (the 3'-nucleotide), and the formation of the first phosphodiester bond were largely unaffected [McClure and Cech, *J.Biol.Chem.* **253**:8949-8956 (1978)]. The dominant effect of rifampicin binding on RNAP activity was a total blockage of synthesis of the second (when transcription was initiated with a nucleoside triphosphate) or third (when transcription was initiated with a nucleoside di- or monophosphate) phosphodiester bond [McClure and Cech, *J.Biol.Chem.* **253**:8949-8956 (1978)]. Since synthesis of the first and second phosphodiester bond can occur in the presence of rifampicin, the

antibiotic does not interfere with substrate binding, catalytic activity, or the intrinsic translocation mechanism of the RNAP. After RNAP has synthesized a long transcript and entered the elongation phase, it becomes totally resistant to rifampicin. These properties led to the proposal that rifampicin inhibits RNAP through a simple steric block of the path of the elongating RNA at the 5'-end [McClure and Cech, *J.Biol.Chem.* **253**:8949-8956 (1978)]. Whether rifampicin directly blocked the path of the RNA, or if blockage was an indirect effect due to a conformational change in the RNAP induced by rifampicin binding, could not be distinguished. It has alternatively been proposed that rifampicin exerts its effect allosterically by decreasing the affinity of the RNAP for short RNA transcripts [Schulz and Zillig, *Nucl.AcidsRes.* **9**:6889-6906 (1981)].

The Rif-RNAP crystal structure explains the results described above and strongly supports the simple steric block mechanism, *see*, atomic coordinates included in Table 2 [McClure and Cech, *J.Biol.Chem.* **253**:8949-8956 (1978)]. Rifampicin directly abuts the base of a loop that comprises the C-terminal part of the β conserved region D (amino acid residues 443-451, shaded red in Figs. 5a-5b), and a cluster of Rif^R mutants, Rif cluster I (Fig. 1), flanks this region. Modeling suggests that this loop, which contains several nearly universally conserved residues, participates in forming the binding site for the base-pair at +1 in the transcription complex [Korzheva *et al.*, *Science* **289**:619-625 (2000)], so effects of rifampicin on the K_m for the initiating substrate are not surprising. However, rifampicin does not directly contact the end of this loop. In addition, conformational changes of the protein in this region are not indicated from the structural data, consistent with the observation that the effect of rifampicin on this region is small.

The principal effect of rifampicin is seen in the context of a model of the transcriptionally active ternary complex [Korzheva *et al.*, *Science* **289**:619-625 (2000)] containing RNAP, DNA template, and RNA transcript (Fig. 6a). In Figure 6, only the RNAP active site Mg^{2+} and the 9-basepair RNA/DNA hybrid (from +1 to -7) from the ternary complex model are shown. The rest of the RNAP and nucleic acids are omitted

for clarity. Also shown is the atomic model of rifampicin as it would be positioned in its binding site on the β subunit.

It can be seen that the two substrate nucleotides, at +1 (green) and -1, are not directly affected by the presence of rifampicin so that RNAP can bind and catalyze the
 5 formation of a phosphodiester bond between the two substrates in the presence of the antibiotic. With a transcript length of 3 nucleotides (nt), however, the 5'-phosphates of the 5'-nucleotide (at -2) sterically clash with rifampicin, and the nucleotides further upstream (-3 to -5) severely clash with rifampicin. At the same time, rifampicin does not interfere with the DNA (grey). Thus, the structure, in combination with the ternary
 10 complex model, explains the biochemical data on the mechanism of rifampicin inhibition, provides strong support for the proposal that rifampicin sterically blocks the path of the elongating RNA transcript at the 5'-end, and indicates that the blockage is a direct consequence of rifampicin binding in its site. The model further suggests why transcripts initiated with nucleoside triphosphates are blocked after the first
 15 phosphodiester bond, while transcripts initiated with nucleoside di- or monophosphates are blocked after the second phosphodiester bond. In the model, the nucleoside monophosphate in the transcript at the -2 position clashes only slightly with rifampicin, while the presence of a 5'-triphosphate at the -2 position would extend into rifampicin.

Core RNAP can bind a pre-formed 'minimal nucleic acid scaffold' of RNA/DNA
 20 oligonucleotides (Fig. 6b, top) to yield functional ternary elongation complexes [Korzheva *et al.*, *Science* **289**:619-625 (2000)]. Order of addition experiments were performed using this system in order to assess whether rifampicin and RNA binding were competitive (Fig. 6b). The DNA component of the scaffold was annealed with varying lengths of RNA transcript, and the effect of rifampicin on the
 25 sequence-dependent extension of RNA by one nucleotide (radioactively-labeled CTP) added before or after the oligonucleotides was assayed at room temperature. In the case of *E. coli* core RNAP in the absence of rifampicin the RNA transcript was extended with nearly equal efficiency regardless of its length within a range of 3-7 nucleotides (Fig. 6b, lanes 11-15). When rifampicin was added prior to the nucleotide

scaffold, the RNAP was unable to extend any of the RNA oligos, regardless of length (lanes 1-5), indicating that rifampicin occupied its site and blocked the extension and/or binding of all of the transcripts. When the scaffold was added prior to rifampicin addition, rifampicin was able to occupy its site and block the extension of the 3-nucleotide transcript (lane 6), but had no effect on the extension of the longer transcripts (lanes 7-10), presumably because rifampicin could not access its binding site due to the presence of the longer RNA transcripts (Fig. 6a). This result is consistent with the early data that rifampicin inhibits the RNA extension from 2 to 3 nucleotides if the 5'-nucleoside is tri-phosphorylated, but inhibits extension from 3 to 4 nucleotides if the 5'-nucleoside is mono- or di-phosphorylated [McClure and Cech, *J.Biol.Chem.* **253**:8949-8956 (1978)] since the synthetic RNA oligos lack 5'-phosphates.

Similar experiments were performed with *Taq* core RNAP (Fig. 6b, lanes 16-30). In the absence of rifampicin, the efficiency of transcript extension was strongly dependent on the transcript length (lanes 26-30). Extension of the shortest transcripts was barely detectable, suggesting that, unlike *E. coli* RNAP, *Taq* core RNAP does not bind and stabilize the short, intrinsically unstable RNA/DNA hybrids. In the presence of rifampicin, a generalized inhibition of transcript extension was observed regardless of the order of addition or of the transcript length (lanes 16-25). These results can be explained by the low binding affinity of *Taq* core RNAP for both rifampicin and for short RNA transcripts compared with *E. coli* core RNAP. The low affinities imply fast off-rates, which would allow equilibrium to be established between the rifampicin and scaffold binding during the time of the assay.

Discussion

The 3.3 Å X-ray crystal structure of *Taq* core RNAP complexed with rifampicin is disclosed herein. Though *Taq* RNAP is less sensitive to rifampicin than *E.coli* rifampicin, at sufficiently high concentrations the antibiotic binds and inhibits the enzyme. Significantly, however, the inhibition of *Taq* RNAP by rifampicin occurs through the same biochemical mechanism as *E. coli* RNAP, and the disposition of the

Rif-site with respect to the active site is identical to *E. coli* RNAP as well as with other prokaryotic RNAPs (Figs. 2a-2d). Therefore, the structural information provided herein is relevant for all bacteria RNAPs.

The relative insensitivity of *Taq* RNAP to rifampicin is likely due to amino acid
 5 substitutions in *Taq* RNAP compared with other, more Rif-sensitive RNAPs. The 12
 residues close enough to interact directly with the rifampicin are identical between *E.*
coli, *Taq*, and *M. tuberculosis* (marked yellow in Fig. 1). Among the 11 secondary
 positions that do not directly interact with rifampicin but likely affect rifampicin
 binding indirectly, 5 are substituted in *Taq* RNAP (amino acid residues 387, 395, 398,
 10 453, and 566; Fig. 1). Three of these positions, 387, 398, and 453, contain amino acids
 that are not dramatically different in overall size from their *E. coli* and *M. tuberculosis*
 counterparts and one would predict that these residues are not the origin of the *Taq*
 RNAP insensitivity to rifampicin. Position 566 is highly conserved among all RNAPs
 as either a lysine or an arginine (the homologous position is an arginine in both *E. coli*
 15 and *M. tuberculosis*) but is a threonine in *Taq* RNAP. This substitution is unlikely to
 be the main determinant of the *Taq* RNAP Rif insensitivity, however, since mutating
Taq Thr566 to an arginine has little effect on the Rif^R of the enzyme when assayed at
 45°C. This leaves position 395, which is highly conserved as a hydrophobic residue
 among all RNAPs. In *E. coli* and *M. tuberculosis* this position is a methionine, but in
 20 *Taq* it is a lysine. *Taq* Lys395 appears to participate in buried salt-bridges with
 Asp124 and Asp133 that may contribute to the thermostability of the protein. This
 non-conservative substitution (lysine for methionine) could affect the local path of the
 polypeptide backbone, and is immediately adjacent to Phe394, the backbone amide and
 carboxyl of which appear to be involved in important interactions with the rifampicin
 25 (Figs. 4a-4b).

All but one of the residues that are close enough to rifampicin to participate in direct
 interactions are known to mutate to strong Rif^R (Figs. 4a-4b). However, additional
 residues could be important for the formation of the Rif binding pocket but not
 revealed as Rif^R mutants if they are necessary for basic RNAP function. As mentioned

above, the four regions of the β subunit that harbor Rif^R mutants are highly conserved among prokaryotes (Fig. 1), but the much weaker homology with archaeobacterial and eukaryotic RNAPs, combined with the fact that so many Rif^R mutations have been discovered, indicate that these regions are not critical to RNAP function *in vivo*.

- 5 Nevertheless, some Rif^R mutations do have profound functional effects [Jin and Gross, *J.Biol.Chem.* **266**:14478-14485 (1991); Landick *et al.*, *Genes Develop.* **4**:1623-1636 (1990)], and *E. coli* strains with Rif^R RNAP have been shown to be at a competitive disadvantage to wild type *E. coli* in the absence of rifampicin [Jin and Gross, *J.Bact.* **171**:5229-5231 (1989)].
- 10 The clinical success of rifampicin proves that the bacterial RNAP is an excellent target for antimicrobials. The structure and available genetic and biochemical data suggest that the design of modified versions of rifampicin to overcome the effects of Rif^R mutations may lead to incremental improvements, though may not lead to a "wonder" drug because of the apparently small functional penalties of mutating this region of the
- 15 RNAP, and the variety of amino acid positions and mutations that result in Rif^R (Fig. 1). In contrast, however, the findings from clinical isolates of Rif^R *M. tuberculosis* are rather encouraging. Thus, although the Rif^R mutations are spread over 15 positions of *rpoB*, 77% of all the mutations isolated involved substitutions at one of only two positions, corresponding to *Taq* amino acid residues 406 and 411. If a third amino acid
- 20 residue is included, *i.e.*, (*Taq* 396) a combined 86% of all the reported mutants are accounted for.

One important conclusion from the present disclosure emerges regarding the inhibitory mechanism of rifampicin, *i.e.*, it is a simple steric block of transcription elongation. Thus, the powerful effects of rifampicin do not stem from the details of its chemical

- 25 structure, and do not involve interference with the catalytic activity of the RNAP, *e.g.*, by mimicking substrates or a transition state of the polymerization reaction. Indeed, such an inhibitor would likely act on features that are highly conserved between prokaryotes and eukaryotes, rendering the inhibitor useless as an antimicrobial agent. Rather, the effects of rifampicin depend only on its ability to bind tightly to a relatively

non-conserved part of the structure, disrupting a critical RNAP function by virtue of its presence. Decades of functional studies [Chamberlin, *Harvey Lectures* 88:1-21 (1993); Korzheva *et al.*, *Cold Spring Harbor Symposia on Quantitative Biology* 63:337-345 (1998); Mustaev *et al.*, *Proc.Nat.Acad.Sci.USA* 91:12036-12040 (1994);
5 and Nudler, *J.Molec.Biol.* 288:1-12 (1999)], and more recent structural evidence [Cramer *et al.*, *Science* 288:640-649 (2000); Korzheva *et al.*, *Science* 289:619-625 (2000); Mooney and Landick, *Cell* 98:687-690(1999); Zhang *et al.*, *Cell* 98:811-824 (1999); U.S. Serial No.09/396,651, Filed September 15, 1999, the contents of which are hereby incorporated by reference in their entireties] indicate that cellular RNAPs
10 operate as complex molecular machines, with extensive interactions with the template DNA, product RNA [Korzheva *et al.*, *Science* 289:619-625 (2000)], and other regulatory molecules. Thus, many additional distinct sites exist where the tight binding of a small molecule (*i.e.*, a novel antibiotic) would disrupt critical features of the functional mechanism of bacterial RNAPs. Such distinct sites can be readily identified
15 through the structural information provided by the present invention.

The present invention is not to be limited in scope by the specific embodiments describe herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and the accompanying figures. Such modifications are intended to fall
20 within the scope of the appended claims.

It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description.

Various publications are cited herein, the disclosures of which are incorporated by
25 reference in their entireties.